

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/62, C07K 16/28, 16/46	A1	(11) International Publication Number: WO 98/23761 (43) International Publication Date: 4 June 1998 (04.06.98)
(21) International Application Number: PCT/US97/19041 (22) International Filing Date: 20 October 1997 (20.10.97) (30) Priority Data: 08/757,205 27 November 1996 (27.11.96) US (71) Applicant: GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US). (72) Inventors: JARDIEU, Paula, M.; 1896 Pacific Avenue #104, San Francisco, CA 94109 (US). PRESTA, Leonard, G.; 1900 Gough Street #206, San Francisco, CA 94109 (US). (74) Agents: LEE, Wendy, M. et al.; Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080-4990 (US).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: HUMANIZED ANTI-CD11a ANTIBODIES (57) Abstract Humanized anti-CD11a antibodies and various uses therefor are disclosed. The humanized anti-CD11a antibody may bind specifically to human CD11a I-domain, have an IC ₅₀ (nM) value of no more than about 1nM for preventing adhesion of Jurkat cells to normal human epidermal keratinocytes expressing ICAM-1, and/or an IC ₅₀ (nM) value of no more than about 1nM in the mixed lymphocyte response assay.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HUMANIZED ANTI-CD11a ANTIBODIES**BACKGROUND OF THE INVENTION****Field of the Invention**

This invention relates generally to humanized anti-CD11a antibodies.

5 Description of Related Art

Lymphocyte function-associated antigen 1 (LFA-1; CD11a/CD18) is involved in leukocyte adhesion during cellular interactions essential for immunologic responses and inflammation (Larson *et al.*, *Immunol. Rev.* 114:181-217 (1990)). LFA-1 is a member of the $\beta 2$ integrin family and consists of a unique α subunit, CD11a, and a β subunit, CD18, common to other $\beta 2$ integrin receptors Mac-1 and p150,95. The ligands of LFA-1 include intercellular adhesion molecule-1, ICAM-1, expressed on leukocytes, endothelium, and dermal fibroblasts (Dustin *et al.*, *J. Immunol.* 137:245-254 (1986)), ICAM-2 expressed on resting endothelium and lymphocytes (de Fougerolles *et al.*, *J. Exp. Med.* 174:253-267 (1991)), and ICAM-3 expressed on monocytes and resting lymphocytes (de Fougerolles *et al.*, *J. Exp. Med.* 179:619-629 (1994)).

Monoclonal antibodies (MAbs) against LFA-1 and the ICAMs have been shown, *in vitro*, to inhibit several T cell-dependent immune functions including T cell activation (Kuypers *et al.*, *Res. Immunol.* 140:461(1989)), T cell-dependent B cell proliferation (Fischer *et al.*, *J. Immunol.* 136:3198-3203 (1986)), target cell lysis (Krensky *et al.*, *J. Immunol.* 131:611-616 (1983)), and adhesion of T cells to vascular endothelium (Lo *et al.*, *J. Immunol.* 143:3325-3329 (1989)). In mice, anti-CD11a MAbs induce tolerance to protein antigens (Tanaka *et al.*, *Eur. J. Immunol.* 25:1555-1558 (1995)) and prolong survival of cardiac (Cavazzana-Calvo *et al.*, *Transplantation* 59:1576-1582 (1995); Nakakura *et al.*, *Transplantation* 55:412-417 (1993)), bone marrow (Cavazzana-Calvo *et al.*, *Transplantation* 59:1576-1582 (1995); van Dijken *et al.*, *Transplantation* 49:882-886 (1990)), corneal (He *et al.*, *Invest. Ophthalmol. Vis. Sci.* 35:3218-3225 (1994)), islet (Nishihara *et al.*, *Transplantation Proc.* 27:372 (1995)) and thyroid (Talento *et al.*, *Transplantation* 55:418-422 (1993)) allografts.

In humans, anti-CD11a MAbs prevent graft failure after bone marrow transplantation (Fischer *et al.*, *Blood* 77:249-256 (1991); Stoppa *et al.*, *Transplant Intl.* 4:3-7 (1991)) and preliminary clinical studies of renal allografts treated prophylactically with anti-CD11a MAb, in addition to corticosteroids and azathioprine, are promising (Hourmant *et al.*, *Transplantation* 58:377-380 (1994)). Current therapies against graft rejection include use of OKT3, a murine anti-human CD3 MAb, and cyclosporin A. OKT3 therapy is effective but has several undesirable side effects; its use results in the release of numerous cytokines including tumor necrosis factor- α , interferon- γ , interleukin-2, and interleukin-6, resulting in fever, chills and gastrointestinal distress (for a review see Parlevliet *et al.*, *Transplant Intl.* 5:234-246 (1992); Dantal *et al.*, *Curr. Opin. Immunol.* 3:740-747 (1991)). Cyclosporin A is effective but also has serious side effects (for a review see Barry, *Drugs*, 44:554-566 (1992)).

35

SUMMARY OF THE INVENTION

The instant invention provides humanized anti-CD11a antibodies. Preferred antibodies bind to the I-domain of human CD11a (*e.g.* to "epitope MHM24" as herein defined) and/or bind CD11a with an affinity of about 1×10^{-8} M or stronger. In preferred embodiments, the antibody has an IC50 (nM) value of no more than about 1nM for preventing adhesion of Jurkat cells to normal human epidermal keratinocytes expressing ICAM-1. Preferred humanized antibodies are those which have an IC50 (nM) value of no more than about

40

1nM in the mixed lymphocyte response (MLR) assay. This IC50 for a humanized antibody in the MLR assay is significantly better than that for murine MAb 25.3, which has been previously tested *in vivo* (Fischer *et al.*, *Blood* 77:249-256 (1991); Stoppa *et al.*, *Transplant Intl.* 4:3-7 (1991); Hourmant *et al.*, *Transplantation* 58:377-380 (1994)).

5 The humanized anti-CD11a antibody may have a heavy chain variable region comprising the amino acid sequence of CDR1 (GYSFTGHWMN; SEQ ID NO:10) and/or CDR2 (MIHPSDSETRYNQKFKD; SEQ ID NO:11) and/or CDR3 (GIYFYGTTFYFDY; SEQ ID NO:12) of humanized antibody MHM24 F(ab)-8 in Fig. 1 and/or a light chain variable region comprising the amino acid sequence of CDR1 (RASKTISKYLA; SEQ ID NO:13) and/or CDR2 (SGSTLQS; SEQ ID NO:14) and/or CDR3 (QQHNEYPLT; SEQ ID NO:15) of humanized antibody MHM24 F(ab)-8 in Fig. 1. In other embodiments, the antibody comprises an amino acid sequence variant of one or more of the CDRs of humanized MHM24 antibody F(ab)-8, which variant comprises one or more amino acid insertion(s) within or adjacent to a CDR residue and/or deletion(s) within or adjacent to a CDR residue and/or substitution(s) of CDR residue(s) (with substitution(s) being the preferred type of amino acid alteration for generating such variants). Such variants will normally have a binding affinity for human CD11a which is no more than about 1×10^{-8} M.

15 In preferred embodiments, the humanized antibody includes a light chain variable region comprising the amino acid sequence of SEQ ID NO:2 and/or a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:5 of humanized antibody MHM24 F(ab)-8 in Fig. 1 and/or amino acid sequence variants thereof.

20 As described herein, it has been possible to reengineer a humanized antibody that bound human CD11a antigen, but not significantly to rhesus CD11a antigen, so as to confer an ability to bind to rhesus CD11a (*i.e.* a "rhesusized" antibody). In this embodiment, the antibody which binds rhesus CD11a may, for example, comprise the CDR2 amino acid sequence in SEQ ID NO:23. The other CDRs may be the same as those for humanized MHM24 antibody F(ab)-8. Thus, the antibody may comprise the amino acid sequence of the "rhesusized" heavy chain in SEQ ID NO:24, optionally combined with a light chain comprising the amino acid sequence in SEQ ID NO:2.

25 Various forms of the antibody are contemplated herein. For example, the anti-CD11a antibody may be a full length antibody (*e.g.* having a human immunoglobulin constant region) or an antibody fragment (*e.g.* a F(ab')₂). Furthermore, the antibody may be labeled with a detectable label, immobilized on a solid phase and/or conjugated with a heterologous compound (such as a cytotoxic agent).

30 Diagnostic and therapeutic uses for the antibody are contemplated. In one diagnostic application, the invention provides a method for determining the presence of CD11a protein comprising exposing a sample suspected of containing the CD11a protein to the anti-CD11a antibody and determining binding of the antibody to the sample. For this use, the invention provides a kit comprising the antibody and instructions for using the antibody to detect the CD11a protein.

35 The invention further provides: isolated nucleic acid encoding the antibody; a vector comprising that nucleic acid, optionally operably linked to control sequences recognized by a host cell transformed with the vector; a host cell comprising that vector; a process for producing the antibody comprising culturing the host cell so that the nucleic acid is expressed and, optionally, recovering the antibody from the host cell culture (*e.g.*

from the host cell culture medium). The invention also provides a composition comprising the humanized anti-CD11a antibody and a pharmaceutically acceptable carrier or diluent. This composition for therapeutic use is sterile and may be lyophilized. The invention further provides a method for treating a mammal suffering from a LFA-1 mediated disorder, comprising administering a pharmaceutically effective amount of the humanized anti-CD11a antibody to the mammal. For such therapeutic uses, other immunosuppressive agents or adhesion molecule antagonists (e.g. another LFA-1 antagonist or a VLA-4 antagonist) may be co-administered to the mammal either before, after, or simultaneously with, the humanized anti-CD11a antibody.

Brief Description of the Drawings

Figure 1A shows the amino acid sequences of murine MHM24 light chain (SEQ ID NO: 1), humanized MHM24 F(ab)-8 light chain (SEQ ID NO:2), human consensus sequences of light chain subgroup κ 1 (hum κ 1) (SEQ ID NO:3).

Figure 1B shows the amino acid sequences of murine MHM24 heavy chain (SEQ ID NO:4), humanized MHM24 F(ab)-8 heavy chain (SEQ ID NO:5), human consensus sequences of heavy chain subgroup III (humIII) (SEQ ID NO:6) and "rhesusized" antibody mutant heavy chain of the Example (SEQ ID NO:24).

In Figs. 1A and 1B, hypervariable regions based on sequence hypervariability (Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) are enclosed within brackets and hypervariable loops based on structure of F(ab)-antigen complexes (Chothia *et al.*, *Nature* 342:8767 (1989)) are in italics. Residue numbering is according to Kabat *et al.*, with insertions shown as a, b, and c.

Figure 2 shows sequences of human CD11a I-domain (SEQ ID NO:7) and rhesus CD11a I-domain (SEQ ID NO:8). β -strands and α -helices are underlined and labeled according to Qu *et al.*, *Proc. Natl. Acad. Sci.* 92:10277-10281 (1995). The rhesus I-domain sequence (rhCD11a) shows only the four differences from human I-domain. The binding epitope for the MHM24 MAbs (SEQ ID NO:9) is shown in bold (Champe *et al.*, *J. Biol. Chem.* 270:1388-1394 (1995)).

Figure 3 depicts inhibition of human Jurkat T-cells to normal human keratinocytes by murine MHM24 (filled circles), chimeric MHM24 (open triangles), humanized MHM24 (HuIgG1) (filled squares), and a human IgG1 isotype control (+). Percent binding measured by fluorescence of labeled Jurkat cells.

Figures 4A-4C show inhibition of binding of rhesus lymphocytes to normal human keratinocytes (Fig. 4A), rhesus lymphocytes to recombinant human ICAM-1 coated on plates (Fig. 4B), and rhesus/human CD11a chimera-transfected 293 cells to normal human keratinocytes (Fig. 4C). Inhibition by rhesus-binding MHM24 (RhIgG1) (filled squares), anti-CD18 MHM23 (filled circles), a human IgG1 isotype control (+) (Figs. 4A and 4C), and a murine IgG1 isotype control (+) (Fig. 4B). Percent binding measured by fluorescence of labeled lymphocytes (Figs. 4A and B) or labeled 293 cells (Fig. 4C).

Figure 5 shows human mixed lymphocyte response assay (MLR) is blocked by murine MHM24 (filled circles), humanized MHM24 (HuIgG1) (filled squares), and a humanized isotype IgG1 control (filled diamond). Percent stimulation index (%SI) is the ratio of the response at a given MAbs concentration to the maximal response with no MAbs present. Data is representative of multiple assays using at least two different stimulator/responder pairs.

Detailed Description of the Preferred Embodiments

I. Definitions

Unless indicated otherwise, the term "CD11a" when used herein refers to the alpha subunit of LFA-1 from any mammal, but preferably from a human. The CD11a may be isolated from a natural source of the molecule or may be produced by synthetic means (e.g., using recombinant DNA technology.) The amino acid sequence for human CD11a is described in EP 362 526B1, for example.

The term "I-domain" of CD11a refers to the region of this molecule delineated in Champe *et al.*, *J. Biol. Chem.* 270:1388-1394 (1995) and/or Qu *et al. Proc. Natl. Acad. Sci.* 92:10277-10281 (1995). The amino acid sequences of human CD11a I-domain (SEQ ID NO:7) and rhesus CD11a I-domain (SEQ ID NO:8) are depicted in Fig. 2 herein.

The term "epitope MHM24" when used herein, unless indicated otherwise, refers to the region in the I-domain of human CD11a to which the MHM24 antibody (see Example below) binds. This epitope comprises the amino acid sequence of SEQ ID NO:9 and, optionally, other amino acid residues of CD11a and/or CD18.

The term "LFA-1-mediated disorder" refers to a pathological state caused by cell adherence interactions involving the LFA-1 receptor on lymphocytes. Examples of such disorders include T cell inflammatory responses such as inflammatory skin diseases including psoriasis; responses associated with inflammatory bowel disease (such as Crohn's disease and ulcerative colitis); adult respiratory distress syndrome; dermatitis; meningitis; encephalitis; uveitis; allergic conditions such as eczema and asthma; conditions involving infiltration of T cells and chronic inflammatory responses; skin hypersensitivity reactions (including poison ivy and poison oak); atherosclerosis; leukocyte adhesion deficiency; autoimmune diseases such as rheumatoid arthritis, systemic lupus erythematosus (SLE), diabetes mellitus, multiple sclerosis, Reynaud's syndrome, autoimmune thyroiditis, experimental autoimmune encephalomyelitis, Sjorgen's syndrome, juvenile onset diabetes, and immune responses associated with delayed hypersensitivity mediated by cytokines and T-lymphocytes typically found in tuberculosis, sarcoidosis, polymyositis, granulomatosis and vasculitis; pernicious anemia; chronic obstructive pulmonary disease (COPD); bronchitis; insulinitis; rhinitis; urticaria; glomerulonephritis; diseases involving leukocyte diapedesis; CNS inflammatory disorder; multiple organ injury syndrome secondary to septicemia or trauma; autoimmune hemolytic anemia; myethemia gravis; antigen-antibody complex mediated diseases; nephrotic syndrome; malignancies (e.g., B-cell malignancies such as chronic lymphocytic leukemia or hairy cell leukemia); all types of transplantations, including graft vs. host or host vs. graft disease; HIV and rhinovirus infection; pulmonary fibrosis; invasion of tumor cells into secondary organs etc.

The term "immunosuppressive agent" as used herein for adjunct therapy refers to substances that act to suppress or mask the immune system of the host into which the graft is being transplanted. This would include substances that suppress cytokine production, downregulate or suppress self-antigen expression, or mask the MHC antigens. Examples of such agents include 2-amino-6-aryl-5-substituted pyrimidines (see U.S. Pat. No. 4,665,077), azathioprine (or cyclophosphamide, if there is an adverse reaction to azathioprine); bromocryptine; glutaraldehyde (which masks the MHC antigens, as described in U.S. Pat. No. 4,120,649); anti-idiotypic antibodies for MHC antigens and MHC fragments; cyclosporin A; steroids such as glucocorticosteroids, e.g., prednisone, methylprednisolone, and dexamethasone; cytokine or cytokine receptor

antagonists including anti-interferon- γ , - β , or - α antibodies; anti-tumor necrosis factor- α antibodies; anti-tumor necrosis factor- β antibodies; anti-interleukin-2 antibodies and anti-IL-2 receptor antibodies; anti-L3T4 antibodies; heterologous anti-lymphocyte globulin; pan-T antibodies, preferably anti-CD3 or anti-CD4/CD4a antibodies; soluble peptide containing a LFA-3 binding domain (WO 90/08187 published 7/26/90); streptokinase; TGF- β ; streptodornase; RNA or DNA from the host; FK506; RS-61443; deoxyspergualin; rapamycin; T-cell receptor (U.S. Pat. No. 5,114,721); T-cell receptor fragments (Offner *et al.*, *Science* 251:430-432 (1991); WO 90/11294; and WO 91/01133); and T cell receptor antibodies (EP 340,109) such as T10B9. These agents are administered at the same time or at separate times from the CD11a antibody, and are used at the same or lesser dosages than as set forth in the art. The preferred adjunct immunosuppressive agent will depend on many factors, including the type of disorder being treated including the type of transplantation being performed, as well as the patient's history, but a general overall preference is that the agent be selected from cyclosporin A, a glucocorticosteroid (most preferably prednisone or methylprednisolone), OKT-3 monoclonal antibody, azathioprine, bromocryptine, heterologous anti-lymphocyte globulin, or a mixture thereof.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, *etc.* Preferably, the mammal is human.

The term "graft" as used herein refers to biological material derived from a donor for transplantation into a recipient. Grafts include such diverse material as, for example, isolated cells such as islet cells and neural-derived cells (*e.g.* schwann cells), tissue such as the amniotic membrane of a newborn, bone marrow, hematopoietic precursor cells, and organs such as skin, heart, liver, spleen, pancreas, thyroid lobe, lung, kidney, tubular organs (*e.g.*, intestine, blood vessels, or esophagus), *etc.* The tubular organs can be used to replace damaged portions of esophagus, blood vessels, or bile duct. The skin grafts can be used not only for burns, but also as a dressing to damaged intestine or to close certain defects such as diaphragmatic hernia. The graft is derived from any mammalian source, including human, whether from cadavers or living donors. Preferably the graft is bone marrow or an organ such as heart and the donor of the graft and the host are matched for HLA class II antigens.

The term "donor" as used herein refers to the mammalian species, dead or alive, from which the graft is derived. Preferably, the donor is human. Human donors are preferably volunteer blood-related donors that are normal on physical examination and of the same major ABO blood group, because crossing major blood group barriers possibly prejudices survival of the allograft. It is, however, possible to transplant, for example, a kidney of a type O donor into an A, B or AB recipient.

The term "transplant" and variations thereof refers to the insertion of a graft into a host, whether the transplantation is syngeneic (where the donor and recipient are genetically identical), allogeneic (where the donor and recipient are of different genetic origins but of the same species), or xenogeneic (where the donor and recipient are from different species). Thus, in a typical scenario, the host is human and the graft is an

isograft, derived from a human of the same or different genetic origins. In another scenario, the graft is derived from a species different from that into which it is transplanted, such as a baboon heart transplanted into a human recipient host, and including animals from phylogenically widely separated species, for example, a pig heart valve, or animal beta islet cells or neuronal cells transplanted into a human host.

5 "Increasing tolerance of a transplanted graft" by a host refers to prolonging the survival of a graft in a host in which it is transplanted, *i.e.*, suppressing the immune system of the host so that it will better tolerate a foreign transplant.

"Intermittent" or "periodic" dosing is a dosing that is continuous for a certain period of time and is at regular intervals that are preferably separated by more than one day.

10 "Selective tolerance" of the disorder refers to a tolerance by the host's immune system for the specific agent causing the disorder, but retaining the ability of the host to reject a second allogeneic or xenogeneic graft. Preferably, the tolerance is such that the immune system is left otherwise intact.

The term "LFA-1 antagonist" refers to a molecule that acts as a competitive inhibitor of the LFA-1 interaction with ICAM-1. Examples of such molecules include antibodies directed against either CD11a (*e.g.*,
15 the humanized anti-CD11a antibodies described herein) or CD18 or both, antibodies to ICAM-1, and other molecules such as peptides (*e.g.*, peptidomimetic antagonists).

The term "VLA-4 antagonist" refers to a molecule that acts as a competitive inhibitor of the VLA-4 interaction with VCAM. Examples of such molecules include antibodies directed against either VLA-4 or VCAM and other molecules (*e.g.*, peptidomimetic antagonists).

20 The term "antibody" is used in the broadest sense and specifically covers monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (*e.g.*, bispecific antibodies), and antibody fragments so long as they exhibit the desired biological activity.

"Antibody fragments" comprise a portion of a full length antibody, generally the antigen binding or variable region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments;
25 diabodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal
30 antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the
35 antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, *Nature* 256:495 (1975), or may be made by recombinant DNA methods (see, *e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature* 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.* 222:581-597 (1991), for example.

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison *et al.*, *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984)).

The term "hypervariable region" when used herein refers to the amino acid residues of an antibody which are responsible for antigen-binding. The hypervariable region comprises amino acid residues from a "complementarity determining region" or "CDR" (*i.e.* residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues from a "hypervariable loop" (*i.e.* residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). "Framework" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined.

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric antibodies which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which hypervariable region residues of the recipient are replaced by hypervariable region residues from a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues which are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature* 321:522-525 (1986); Reichmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992).

"Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Generally, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds. Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy chain variable domain (V_H) connected to a light chain variable domain (V_L) in the same polypeptide chain ($V_H - V_L$). By using a linker that is too short to allow pairing between the two

domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993).

The expression "linear antibodies" when used throughout this application refers to the antibodies described in Zapata *et al.* *Protein Eng.* 8(10):1057-1062 (1995). Briefly, these antibodies comprise a pair of tandem Fd segments ($V_H-C_H1-V_H-C_H1$) which form a pair of antigen binding regions. Linear antibodies can be bispecific or monospecific.

An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

The term "epitope tagged" when used herein refers to the anti-CD11a antibody fused to an "epitope tag". The epitope tag polypeptide has enough residues to provide an epitope against which an antibody thereagainst can be made, yet is short enough such that it does not interfere with activity of the CD11a antibody. The epitope tag preferably is sufficiently unique so that the antibody thereagainst does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least 6 amino acid residues and usually between about 8-50 amino acid residues (preferably between about 9-30 residues). Examples include the flu HA tag polypeptide and its antibody 12CA5 (Field *et al.* *Mol. Cell. Biol.* 8:2159-2165 (1988)); the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto (Evan *et al.*, *Mol. Cell. Biol.* 5(12):3610-3616 (1985)); and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody (Paborsky *et al.*, *Protein Engineering* 3(6):547-553 (1990)). In certain embodiments, the epitope tag is a "salvage receptor binding epitope".

As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (*e.g.*, ¹³¹I, ¹²⁵I, ⁹⁰Y and ¹⁸⁶Re), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant or animal origin, or fragments thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include Adriamycin, Doxorubicin, 5-Fluorouracil, Cytosine arabinoside ("Ara-C"), Cyclophosphamide, Thiotepe, Taxotere (docetaxel), Busulfan, Cytosine, Taxol, Methotrexate, Cisplatin, Melphalan, Vinblastine, Bleomycin, Etoposide, Ifosfamide, Mitomycin C, Mitoxantrone, Vincristine,

Vinorelbine, Carboplatin, Teniposide, Daunomycin, Carminomycin, Aminopterin, Dactinomycin, Mitomycins, Esperamicins (see U.S. Pat. No. 4,675,187), Melphalan and other related nitrogen mustards.

The term "prodrug" as used in this application refers to a precursor or derivative form of a pharmaceutically active substance that is less cytotoxic to tumor cells compared to the parent drug and is capable of being enzymatically activated or converted into the more active parent form. See, e.g., Wilman, "Prodrugs in Cancer Chemotherapy" *Biochemical Society Transactions*, 14, pp. 375-382, 615th Meeting Belfast (1986) and Stella *et al.*, "Prodrugs: A Chemical Approach to Targeted Drug Delivery," *Directed Drug Delivery*, Borchardt *et al.*, (ed.), pp. 247-267, Humana Press (1985). The prodrugs of this invention include, but are not limited to, phosphate-containing prodrugs, thiophosphate-containing prodrugs, sulfate-containing prodrugs, peptide-containing prodrugs, D-amino acid-modified prodrugs, glycosylated prodrugs, β -lactam-containing prodrugs, optionally substituted phenoxyacetamide-containing prodrugs or optionally substituted phenylacetamide-containing prodrugs, 5-fluorocytosine and other 5-fluorouridine prodrugs which can be converted into the more active cytotoxic free drug. Examples of cytotoxic drugs that can be derivatized into a prodrug form for use in this invention include, but are not limited to, those chemotherapeutic agents described above.

The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody. The label may itself be detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

By "solid phase" is meant a non-aqueous matrix to which the antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g. controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g. an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Patent No. 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as the anti-CD11a antibodies disclosed herein and, optionally, a chemotherapeutic agent) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

An "isolated" nucleic acid molecule is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the antibody nucleic acid. An isolated nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from the nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule includes a nucleic acid molecule contained in cells that ordinarily express the antibody where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The expression "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for

prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

As used herein, the expressions "cell," "cell line," and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Mutant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. Where distinct designations are intended, it will be clear from the context.

II. Modes for Carrying out the Invention

A. Antibody Preparation

A method for humanizing a nonhuman CD11a antibody is described in the Example below. In order to humanize an anti-CD11a antibody, the nonhuman antibody starting material is prepared. Exemplary techniques for generating such antibodies will be described in the following sections.

(i) Antigen preparation.

The CD11a antigen to be used for production of antibodies may be, *e.g.*, a soluble form of the extracellular domain of CD11a or other fragment of CD11a (*e.g.* a CD11a fragment comprising the "MHM24 epitope", such as CD11a I-domain fragment). Alternatively, cells expressing CD11a at their cell surface can be used to generate antibodies. Such cells can be transformed to express CD11a and, optionally, CD18 or may be other naturally occurring cells (*e.g.* human lymphoblastoid cells, see Hildreth *et al. Eur. J. Immunol.* 13:202-208 (1983)) or Jurkat cells (see Example below). Other forms of CD11a useful for generating antibodies will be apparent to those skilled in the art.

(ii) Polyclonal antibodies

Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the relevant antigen to a protein that is immunogenic in the species to be immunized, *e.g.*, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl groups.

Animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, e.g., 100 µg or 5 µg of the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitably used to enhance the immune response.

10 (iii) *Monoclonal antibodies*

Monoclonal antibodies may be made using the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster or macaque monkey, is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)).

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOP-21 and M.C.-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson *et al.*, *Anal. Biochem.*, 107:220 (1980).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may
5 be grown *in vivo* as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional
10 procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the monoclonal antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the
15 recombinant host cells. Recombinant production of antibodies will be described in more detail below.

(iv) *Humanization and amino acid sequence variants*

The Example below describes a procedure for humanization of an anti-CD11a antibody. In certain embodiments, it may be desirable to generate amino acid sequence variants of the humanized antibody, particularly where these improve the binding affinity or other biological properties of the humanized antibody.

20 Amino acid sequence variants of humanized anti-CD11a antibody are prepared by introducing appropriate nucleotide changes into the humanized anti-CD11a antibody DNA, or by peptide synthesis. Such variants include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences shown for the humanized anti-CD11a F(ab)-8 (e.g. as in SEQ ID NO's 2 & 5). Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the
25 final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the humanized anti-CD11a antibody, such as changing the number or position of glycosylation sites.

A useful method for identification of certain residues or regions of the humanized anti-CD11a antibody polypeptide that are preferred locations for mutagenesis is called "alanine scanning mutagenesis," as
30 described by Cunningham and Wells *Science*, 244:1081-1085 (1989). Here, a residue or group of target residues are identified (e.g., charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with CD11a antigen. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site
35 for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* need not be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed humanized anti-CD11a antibody variants are screened for the desired activity.

Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include humanized anti-CD11a antibody with an N-terminal methionyl residue or the antibody fused to an epitope tag. Other insertional variants of the humanized anti-CD11a antibody molecule include the fusion to the N- or C-terminus of humanized anti-CD11a antibody of an enzyme or a polypeptide which increases the serum half-life of the antibody (see below).

Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the humanized anti-CD11a antibody molecule removed and a different residue inserted in its place. The sites of greatest interest for substitutional mutagenesis include the hypervariable loops, but FR alterations are also contemplated. Table IV in the Example below provides guidance as to hypervariable region residues which can be altered. Hypervariable region residues or FR residues involved in antigen binding are generally substituted in a relatively conservative manner. Such conservative substitutions are shown in Table I under the heading of "preferred substitutions". If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions" in Table I, or as further described below in reference to amino acid classes, are introduced and the products screened.

Table I

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	glu
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro; ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr

Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

5 Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- 10 (1) hydrophobic: norleucine, met, ala, val, leu, ile;
 (2) neutral hydrophilic: cys, ser, thr;
 (3) acidic: asp, glu;
 (4) basic: asn, gln, his, lys, arg;
 (5) residues that influence chain orientation: gly, pro; and
 15 (6) aromatic: trp, tyr, phe.

 Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Any cysteine residue not involved in maintaining the proper conformation of the humanized anti-CD11a antibody also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to
 20 improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

 Another type of amino acid variant of the antibody alters the original glycosylation pattern of the antibody. By altering is meant deleting one or more carbohydrate moieties found in the antibody, and/or adding one or more glycosylation sites that are not present in the antibody.

 Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment
 25 of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acylgalactosamine, galactose, or xylose to a hydroxyamino
 30 acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

 Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

35 Nucleic acid molecules encoding amino acid sequence variants of humanized anti-CD11a antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by

oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of humanized anti-CD11a antibody.

Ordinarily, amino acid sequence variants of the humanized anti-CD11a antibody will have an amino acid sequence having at least 75% amino acid sequence identity with the original humanized antibody amino acid sequences of either the heavy or the light chain (e.g. as in SEQ ID NO:2 or 5), more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, and most preferably at least 95%. Identity or homology with respect to this sequence is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the humanized anti-CD11a residues, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions (as defined in Table I above) as part of the sequence identity. None of N-terminal, C-terminal, or internal extensions, deletions, or insertions into the antibody sequence shall be construed as affecting sequence identity or homology.

(v) *Screening for biological properties*

Antibodies having the characteristics identified herein as being desirable in a humanized anti-CD11a antibody are screened for.

To screen for antibodies which bind to the epitope on CD11a bound by an antibody of interest (e.g., those which block binding of the MHM24 antibody to CD11a), a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Alternatively, epitope mapping, e.g. as described in Champe *et al.*, *J. Biol. Chem.* 270:1388-1394 (1995), can be performed to determine whether the antibody binds an epitope of interest.

Antibody affinities (e.g. for human CD11a or rhesus CD11a) may be determined by saturation binding using either peripheral blood mononuclear cells or rhesus leukocytes as described in the Example below. According to this method for determining antibody affinity, lymphocytes or rhesus leukocytes are added to the plates in a volume of 170 μ l per well and plates are incubated for 2 hr at room. After incubation, cells are harvested and washed 10 times. Samples are then counted. Data is transformed from counts per minute to nanomolarity and four-parameter curve-fitting of saturation plots (bound versus total) are then performed to determine K_d (app) values. Preferred humanized antibodies are those which bind human CD11a with a K_d value of no more than about 1×10^{-7} ; preferably no more than about 1×10^{-8} ; more preferably no more than about 1×10^{-9} ; and most preferably no more than about 2×10^{-10} .

It is also desirable to select humanized antibodies which have beneficial anti-adhesion properties in the "keratinocyte monolayer adhesion assay". Preferred antibodies are those which have an IC_{50} (nM) value of no more than about 250nM; preferably no more than about 100nM; more preferably no more than about 1nM and most preferably no more than about 0.5nM for preventing adhesion of Jurkat cells to normal human epidermal keratinocytes expressing ICAM-1. According to this assay, normal human epidermal keratinocytes are removed from culture flasks and resuspended in lymphocyte assay medium at a concentration of 5×10^5 viable cells/ml. Aliquots of 0.1 ml/well are then cultured overnight in flat-bottom 96-well plates; appropriate wells are stimulated by addition of interferon-gamma at 100 units/well. Jurkat clone E6-1 cells are labeled, washed, resuspended to 1×10^6 cells/ml, and incubated with 2-fold serial dilutions starting at 500ng/ml antibody at 4°C for 30 min. After removal of medium from the keratinocyte monolayer, 0.1 ml/well of labeled

cells are added and incubated at 37°C for 1 h. The wells are washed to remove non-attached cells and fluorescence is measured.

Desirable humanized anti-CD11a antibodies are those which have an IC₅₀ (nM) value of no more than about 100nM; preferably no more than about 50nM; more preferably no more than about 5nM and most preferably no more than about 1nM in the mixed lymphocyte response (MLR) assay, using human lymphocytes. For both human and rhesus MLR, peripheral blood lymphocytes from two unrelated donors are isolated from whole, heparinized blood and are resuspended to a concentration of 3×10^6 cells/ml in RPMI 1640 (GIBCO) with additives as described in the Example below. The stimulator cells are made unresponsive by irradiation. Responder cells at a concentration of 1.5×10^5 cells per well are co-cultured with an equal number of stimulator cells in 96-well, flat-bottom plates. Two-fold serial dilutions of antibody starting at a concentration of 10nM are added to the cultures to give a total volume of 200 µl/well. The cultures are incubated at 37°C in 5% CO₂ for 5 days and then pulsed with 1 µCi/well of [³H]thymidine for 16 h and [³H]thymidine incorporation is measured.

(vi) *Antibody fragments*

In certain embodiments, the humanized CD11a antibody is an antibody fragment. Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992) and Brennan *et al.*, *Science* 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. For example, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, *Bio/Technology* 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner.

(vii) *Multispecific antibodies*

In some embodiments, it may be desirable to generate multispecific (e.g. bispecific) humanized CD11a antibodies having binding specificities for at least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of the CD11a protein. Alternatively, an anti-CD11a arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g., CD2 or CD3), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the CD11a-expressing cell. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express CD11a. These antibodies possess an CD11a-binding arm and an arm which binds the cytotoxic agent (e.g., saporin, anti-interferon-α, vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g., F(ab')₂ bispecific antibodies).

According to another approach for making bispecific antibodies, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g., tyrosine or tryptophan). Compensatory "cavities"

of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g., alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers. See WO96/27011 published September 6, 1996.

5 Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

Techniques for generating bispecific antibodies from antibody fragments have also been described
10 in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science* 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate $F(ab')_2$ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then
15 reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby *et al.*, *J. Exp. Med.* 175:217-225 (1992) describe
20 the production of a fully humanized bispecific antibody $F(ab')_2$ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the HER2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant
25 cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, *J. Immunol.* 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody"
30 technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two
35 antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, *J. Immunol.* 152:5368 (1994). Alternatively, the bispecific antibody may be a "linear antibody" produced as described in Zapata *et al. Protein Eng.* 8(10):1057-1062 (1995).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.*, *J. Immunol.* 147:60 (1991).

(viii) *Other modifications*

Other modifications of the humanized anti-CD11a antibody are contemplated. For example, it may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance the effectiveness of the antibody in treating cancer, for example. For example cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, *J. Exp Med.* 176:1191-1195 (1992) and Shopes, B. *J. Immunol.* 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.*, *Cancer Research* 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.*, *Anti-Cancer Drug Design* 3:219-230 (1989).

The invention also pertains to immunoconjugates comprising the antibody described herein conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated anti-CD11a antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y and ^{186}Re .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl)hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, *Science* 238:1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (*e.g.*, avidin) which is conjugated to a cytotoxic agent (*e.g.*, a radio nuclide).

The anti-CD11a antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, *Proc. Natl. Acad. Sci. USA* 82:3688 (1985); Hwang *et al.*, *Proc. Natl. Acad. Sci. USA* 77:4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.*, *J. Biol. Chem.* 257:286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon *et al.*, *J. National Cancer Inst.* 81(19):1484 (1989)

The antibody of the present invention may also be used in ADEPT by conjugating the antibody to a prodrug-activating enzyme which converts a prodrug (*e.g.*, a peptidyl chemotherapeutic agent, see WO81/01145) to an active anti-cancer drug. See, for example, WO 88/07378 and U.S. Patent No. 4,975,278.

The enzyme component of the immunoconjugate useful for ADEPT includes any enzyme capable of acting on a prodrug in such a way so as to convert it into its more active, cytotoxic form.

Enzymes that are useful in the method of this invention include, but are not limited to, alkaline phosphatase useful for converting phosphate-containing prodrugs into free drugs; arylsulfatase useful for converting sulfate-containing prodrugs into free drugs; cytosine deaminase useful for converting non-toxic 5-fluorocytosine into the anti-cancer drug, 5-fluorouracil; proteases, such as serratia protease, thermolysin, subtilisin, carboxypeptidases and cathepsins (such as cathepsins B and L), that are useful for converting peptide-containing prodrugs into free drugs; D-alanylcarboxypeptidases, useful for converting prodrugs that contain D-amino acid substituents; carbohydrate-cleaving enzymes such as β -galactosidase and neuraminidase useful for converting glycosylated prodrugs into free drugs; β -lactamase useful for converting drugs derivatized with β -lactams into free drugs; and penicillin amidases, such as penicillin V amidase or penicillin G amidase, useful for converting drugs derivatized at their amine nitrogens with phenoxyacetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies with enzymatic activity, also known in the art as "abzymes", can be used to convert the prodrugs of the invention into free active drugs (see, *e.g.*, Massey, *Nature* 328:457-458 (1987)). Antibody-abzyme conjugates can be prepared as described herein for delivery of the abzyme to a tumor cell population.

The enzymes of this invention can be covalently bound to the anti-CD11a antibodies by techniques well known in the art such as the use of the heterobifunctional crosslinking reagents discussed above. Alternatively, fusion proteins comprising at least the antigen binding region of an antibody of the invention linked to at least a functionally active portion of an enzyme of the invention can be constructed using recombinant DNA techniques well known in the art (see, *e.g.*, Neuberger *et al.*, *Nature* 312:604-608 (1984)).

In certain embodiments of the invention, it may be desirable to use an antibody fragment, rather than an intact antibody, to increase tumor penetration, for example. In this case, it may be desirable to modify the antibody fragment in order to increase its serum half life. This may be achieved, for example, by incorporation

of a salvage receptor binding epitope into the antibody fragment (*e.g.*, by mutation of the appropriate region in the antibody fragment or by incorporating the epitope into a peptide tag that is then fused to the antibody fragment at either end or in the middle, *e.g.*, by DNA or peptide synthesis). See WO96/32478 published October 17, 1996.

- 5 The salvage receptor binding epitope generally constitutes a region wherein any one or more amino acid residues from one or two loops of a Fc domain are transferred to an analogous position of the antibody fragment. Even more preferably, three or more residues from one or two loops of the Fc domain are transferred. Still more preferred, the epitope is taken from the CH2 domain of the Fc region (*e.g.*, of an IgG) and transferred to the CH1, CH3, or V_H region, or more than one such region, of the antibody. Alternatively,
- 10 the epitope is taken from the CH2 domain of the Fc region and transferred to the C_L region or V_L region, or both, of the antibody fragment.

- In one most preferred embodiment, the salvage receptor binding epitope comprises the sequence (5' to 3'): PKNSSMISNTP (SEQ ID NO:16), and optionally further comprises a sequence selected from the group consisting of HQSLGTQ (SEQ ID NO:17), HQNLSDGK (SEQ ID NO:18), HQNISDGK (SEQ ID NO:19),
- 15 or VISSHLGQ (SEQ ID NO:20), particularly where the antibody fragment is a Fab or F(ab')₂. In another most preferred embodiment, the salvage receptor binding epitope is a polypeptide containing the sequence(s) (5' to 3'): HQNLSDGK (SEQ ID NO:18), HQNISDGK (SEQ ID NO:19), or VISSHLGQ (SEQ ID NO:20) and the sequence: PKNSSMISNTP (SEQ ID NO:16).

- Covalent modifications of the humanized CD11a antibody are also included within the scope of this
- 20 invention. They may be made by chemical synthesis or by enzymatic or chemical cleavage of the antibody, if applicable. Other types of covalent modifications of the antibody are introduced into the molecule by reacting targeted amino acid residues of the antibody with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

- Cysteiny residues most commonly are reacted with α -haloacetates (and corresponding amines), such
- 25 as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteiny residues also are derivatized by reaction with bromotrifluoroacetone, α -bromo- β -(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

- Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this
- 30 agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0.

- Lysiny and amino-terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysiny residues. Other suitable reagents for derivatizing α -amino-containing residues include imidoesters such as methyl picolinimide,
- 35 pyridoxal phosphate, pyridoxal, chloroborohydride, trinitrobenzenesulfonic acid, O-methylisourea, 2,4-pentanedione, and transaminase-catalyzed reaction with glyoxylate.

Arginy residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pK_a of the guanidine

functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane.

5 Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using ^{125}I or ^{131}I to prepare labeled proteins for use in radioimmunoassay.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides (R-N=C=N-R'), where R and R' are different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl
10 residues are converted to asparaginy and glutaminy residues by reaction with ammonium ions.

Glutaminy and asparaginy residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. These residues are deamidated under neutral or basic conditions. The deamidated form of these residues falls within the scope of this invention.

15 Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains (T.E. Creighton, *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification involves chemically or enzymatically coupling glycosides to
20 the antibody. These procedures are advantageous in that they do not require production of the antibody in a host cell that has glycosylation capabilities for N- or O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of
25 glutamine. These methods are described in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, *CRC Crit. Rev. Biochem.*, pp. 259-306 (1981).

Removal of any carbohydrate moieties present on the antibody may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the antibody to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or
30 all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the antibody intact. Chemical deglycosylation is described by Hakimuddin, *et al. Arch. Biochem. Biophys.* 259:52 (1987) and by Edge *et al. Anal. Biochem.*, 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on antibodies can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura *et al. Meth. Enzymol.* 138:350 (1987).

35 Another type of covalent modification of the antibody comprises linking the antibody to one of a variety of nonproteinaceous polymers, *e.g.*, polyethylene glycol, polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

B. Vectors, Host Cells and Recombinant Methods

The invention also provides isolated nucleic acid encoding the humanized anti-CD11a antibody, vectors and host cells comprising the nucleic acid, and recombinant techniques for the production of the antibody.

5 For recombinant production of the antibody, the nucleic acid encoding it is isolated and inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. DNA encoding the monoclonal antibody is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody). Many vectors are available. The vector components generally include, but are not limited to, 10 one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

(i) *Signal sequence component*

The anti-CD11a antibody of this invention may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which is preferably a signal sequence or other 15 polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. The heterologous signal sequence selected preferably is one that is recognized and processed (*i.e.*, cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the native anti-CD11a antibody signal sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. 20 For yeast secretion the native signal sequence may be substituted by, *e.g.*, the yeast invertase leader, α factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders), or acid phosphatase leader, the *C. albicans* glucoamylase leader, or the signal described in WO 90/13646. In mammalian cell expression, mammalian signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, are available.

25 The DNA for such precursor region is ligated in reading frame to DNA encoding the anti-CD11a antibody.

(ii) *Origin of replication component*

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the 30 vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed 35 for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

(iii) *Selection gene component*

Expression and cloning vectors may contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin,

neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, *e.g.*, the gene encoding D-alanine racemase for *Bacilli*.

One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin, mycophenolic acid and hygromycin.

Another example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the anti-CD11a antibody nucleic acid, such as DHFR, thymidine kinase, metallothionein-I and -II, preferably primate metallothionein genes, adenosine deaminase, ornithine decarboxylase, *etc.*

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity.

Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding anti-CD11a antibody, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH) can be selected by cell growth in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, *e.g.*, kanamycin, neomycin, or G418. See U.S. Patent No. 4,965,199.

A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 (Stinchcomb *et al.*, *Nature*, 282:39 (1979)). The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1. Jones, *Genetics*, 85:12 (1977). The presence of the *trp1* lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, *Leu2*-deficient yeast strains (ATCC 20,622 or 38,626) are complemented by known plasmids bearing the *Leu2* gene.

In addition, vectors derived from the 1.6 μ m circular plasmid pKD1 can be used for transformation of *Kluyveromyces* yeasts. Alternatively, an expression system for large-scale production of recombinant calf chymosin was reported for *K. lactis*. Van den Berg, *Bio/Technology*, 8:135 (1990). Stable multi-copy expression vectors for secretion of mature recombinant human serum albumin by industrial strains of *Kluyveromyces* have also been disclosed. Fleer *et al.*, *Bio/Technology*, 9:968-975 (1991).

(iv) Promoter component

Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the anti-CD11a antibody nucleic acid. Promoters suitable for use with prokaryotic hosts include the *phoA* promoter, β -lactamase and lactose promoter systems, alkaline phosphatase, a tryptophan (*trp*) promoter system, and hybrid promoters such as the *tac* promoter. However, other known bacterial promoters are suitable. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the anti-CD11a antibody.

Promoter sequences are known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence

found 70 to 80 bases upstream from the start of transcription of many genes is a CNCAAT region where N may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

5 Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

10 Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657. Yeast enhancers also are
15 advantageously used with yeast promoters.

 Anti-CD11a antibody transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, *e.g.*, the actin
20 promoter or an immunoglobulin promoter, from heat-shock promoters, provided such promoters are compatible with the host cell systems.

 The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment. A system for expressing DNA
25 in mammalian hosts using the bovine papilloma virus as a vector is disclosed in U.S. Patent No. 4,419,446. A modification of this system is described in U.S. Patent No. 4,601,978. See also Reyes *et al.*, *Nature* 297:598-601 (1982) on expression of human β -interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus. Alternatively, the rous sarcoma virus long terminal repeat can be used as the promoter.

30 (v) *Enhancer element component*

 Transcription of a DNA encoding the anti-CD11a antibody of this invention by higher eukaryotes is often increased by inserting an enhancer sequence into the vector. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the
35 replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, *Nature* 297:17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the anti-CD11a antibody-encoding sequence, but is preferably located at a site 5' from the promoter.

(vi) *Transcription termination component*

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding anti-CD11a antibody. One useful transcription termination component is the bovine growth hormone polyadenylation region. See WO94/11026 and the expression vector disclosed therein.

(vii) *Selection and transformation of host cells*

Suitable host cells for cloning or expressing the DNA in the vectors herein are the prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes for this purpose include eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* X1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for anti-CD11a antibody-encoding vectors. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe*; *Kluyveromyces* hosts such as, e.g., *K. lactis*, *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilarum* (ATCC 36,906), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa*; *Schwanniomyces* such as *Schwanniomyces occidentalis*; and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

Suitable host cells for the expression of glycosylated anti-CD11a antibody are derived from multicellular organisms. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been identified. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can also be utilized as hosts.

However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey

kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, *J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub *et al.*, *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather *et al.*, *Annals N.Y. Acad. Sci.* 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

Host cells are transformed with the above-described expression or cloning vectors for anti-CD11a antibody production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

(viii) *Culturing the host cells*

The host cells used to produce the anti-CD11a antibody of this invention may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ((MEM), (Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham *et al.*, *Meth. Enz.* 58:44 (1979), Barnes *et al.*, *Anal. Biochem.* 102:255 (1980), U.S. Pat. Nos. 4,767,704; 4,657,866; 4,927,762; 4,560,655; or 5,122,469; WO 90/03430; WO 87/00195; or U.S. Patent Re. 30,985 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as GENTAMYCINTM drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

(ix) *Purification of anti-CD11a antibody*

When using recombinant techniques, the antibody can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the antibody is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, is removed, for example, by centrifugation or ultrafiltration. Carter *et al.*, *Bio/Technology* 10:163-167 (1992) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonyl fluoride (PMSF) over about 30 min. Cell debris can be removed by centrifugation. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

The antibody composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the antibody. Protein A can be used to purify antibodies that are based on human $\gamma 1$, $\gamma 2$, or $\gamma 4$ heavy chains (Lindmark *et al.*, *J. Immunol. Meth.* 62:1-13 (1983)). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss *et al.*, *EMBO J.* 5:15671575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a C_H3 domain, the Bakerbond ABXTMresin (J. T. Baker, Phillipsburg, NJ) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSETM chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

Following any preliminary purification step(s), the mixture comprising the antibody of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, preferably performed at low salt concentrations (*e.g.*, from about 0-0.25M salt).

C. Pharmaceutical Formulations

Therapeutic formulations of the antibody are prepared for storage by mixing the antibody having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as TWEENTM, PLURONICSTM or polyethylene glycol (PEG).

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide an immunosuppressive agent.¹ Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsule prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsule and

poly-(methylmethacrylate) microcapsule, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980).

5 The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsule. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate); or poly(vinylalcohol)), polylactides
10 (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the Lupron Depot™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated
15 antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions,
20 controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

D. Non-therapeutic Uses for the Antibody

The antibodies of the invention may be used as affinity purification agents. In this process, the antibodies are immobilized on a solid phase such as Sephadex resin or filter paper, using methods well known
25 in the art. The immobilized antibody is contacted with a sample containing the CD11a protein (or fragment thereof) to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the CD11a protein, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent, such as glycine buffer, pH 5.0, that will release the CD11a protein from the antibody.

30 Anti-CD11a antibodies may also be useful in diagnostic assays for CD11a protein, e.g., detecting its expression in specific cells, tissues, or serum.

For diagnostic applications, the antibody typically will be labeled with a detectable moiety. Numerous labels are available which can be generally grouped into the following categories:

(a) Radioisotopes, such as ^{35}S , ^{14}C , ^{125}I , ^3H , and ^{131}I . The antibody can be labeled with the
35 radioisotope using the techniques described in *Current Protocols in Immunology*, Volumes 1 and 2, Coligen *et al.*, Ed. Wiley-Interscience, New York, New York, Pubs. (1991) for example and radioactivity can be measured using scintillation counting.

(b) Fluorescent labels such as rare earth chelates (europium chelates) or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, Lissamine, phycoerythrin and Texas Red are available. The fluorescent

labels can be conjugated to the antibody using the techniques disclosed in *Current Protocols in Immunology*, *supra*, for example. Fluorescence can be quantified using a fluorimeter.

- (c) Various enzyme-substrate labels are available and U.S. Patent No. 4,275,149 provides a review of some of these. The enzyme generally catalyzes a chemical alteration of the chromogenic substrate which can be measured using various techniques. For example, the enzyme may catalyze a color change in a substrate, which can be measured spectrophotometrically. Alternatively, the enzyme may alter the fluorescence or chemiluminescence of the substrate. Techniques for quantifying a change in fluorescence are described above. The chemiluminescent substrate becomes electronically excited by a chemical reaction and may then emit light which can be measured (using a chemiluminometer, for example) or donates energy to a fluorescent acceptor.
- Examples of enzymatic labels include luciferases (e.g., firefly luciferase and bacterial luciferase; U.S. Patent No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, malate dehydrogenase, urease, peroxidase such as horseradish peroxidase (HRPO), alkaline phosphatase, β -galactosidase, glucoamylase, lysozyme, saccharide oxidases (e.g., glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase), heterocyclic oxidases (such as uricase and xanthine oxidase), lactoperoxidase, microperoxidase, and the like. Techniques for conjugating enzymes to antibodies are described in O'Sullivan *et al.*, *Methods for the Preparation of Enzyme-Antibody Conjugates for use in Enzyme Immunoassay*, in *Methods in Enzym.* (ed J. Langone & H. Van Vunakis), Academic press, New York, 73:147-166 (1981).

Examples of enzyme-substrate combinations include, for example:

- (i) Horseradish peroxidase (HRPO) with hydrogen peroxidase as a substrate, wherein the hydrogen peroxidase oxidizes a dye precursor (e.g., orthophenylene diamine (OPD) or 3,3',5,5'-tetramethyl benzidine hydrochloride (TMB));
- (ii) alkaline phosphatase (AP) with para-Nitrophenyl phosphate as chromogenic substrate; and
- (iii) β -D-galactosidase (β -D-Gal) with a chromogenic substrate (e.g., p-nitrophenyl- β -D-galactosidase) or fluorogenic substrate 4-methylumbelliferyl- β -D-galactosidase.

- Numerous other enzyme-substrate combinations are available to those skilled in the art. For a general review of these, see U.S. Patent Nos. 4,275,149 and 4,318,980.

Sometimes, the label is indirectly conjugated with the antibody. The skilled artisan will be aware of various techniques for achieving this. For example, the antibody can be conjugated with biotin and any of the three broad categories of labels mentioned above can be conjugated with avidin, or *vice versa*. Biotin binds selectively to avidin and thus, the label can be conjugated with the antibody in this indirect manner. Alternatively, to achieve indirect conjugation of the label with the antibody, the antibody is conjugated with a small hapten (e.g., digoxin) and one of the different types of labels mentioned above is conjugated with an anti-hapten antibody (e.g., anti-digoxin antibody). Thus, indirect conjugation of the label with the antibody can be achieved.

- In another embodiment of the invention, the anti-CD11a antibody need not be labeled, and the presence thereof can be detected using a labeled antibody which binds to the CD11a antibody.

The antibodies of the present invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, *Monoclonal Antibodies: A Manual of Techniques*, pp.147-158 (CRC Press, Inc. 1987).

Competitive binding assays rely on the ability of a labeled standard to compete with the test sample analyte for binding with a limited amount of antibody. The amount of CD11a protein in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies generally are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte which remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex. See, e.g., US Pat No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

For immunohistochemistry, the tumor sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin, for example.

The antibodies may also be used for *in vivo* diagnostic assays. Generally, the antibody is labeled with a radio nuclide (such as ^{111}In , ^{99}Tc , ^{14}C , ^{131}I , ^{125}I , ^3H , ^{32}P or ^{35}S) so that the tumor can be localized using immunoscintigraphy.

E. Diagnostic Kits

As a matter of convenience, the antibody of the present invention can be provided in a kit, *i.e.*, a packaged combination of reagents in predetermined amounts with instructions for performing the diagnostic assay. Where the antibody is labeled with an enzyme, the kit will include substrates and cofactors required by the enzyme (*e.g.*, a substrate precursor which provides the detectable chromophore or fluorophore). In addition, other additives may be included such as stabilizers, buffers (*e.g.*, a block buffer or lysis buffer) and the like. The relative amounts of the various reagents may be varied widely to provide for concentrations in solution of the reagents which substantially optimize the sensitivity of the assay. Particularly, the reagents may be provided as dry powders, usually lyophilized, including excipients which on dissolution will provide a reagent solution having the appropriate concentration.

F. Therapeutic Uses for the Antibody

It is contemplated that the anti-CD11a antibody of the present invention may be used to treat the various LFA-1 mediated disorders as described herein.

The anti-CD11a antibody is administered by any suitable means, including parenteral, subcutaneous, intraperitoneal, intrapulmonary, and intranasal, and, if desired for local immunosuppressive treatment, intralesional administration (including perfusing or otherwise contacting the graft with the antibody before transplantation). Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. In addition, the anti-CD11a antibody is suitably administered by pulse infusion, particularly with declining doses of the antibody. Preferably the dosing is given by injections, most preferably intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic.

For the prevention or treatment of disease, the appropriate dosage of antibody will depend on the type of disease to be treated, as defined above, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

Depending on the type and severity of the disease, about 1 µg/kg to 15 mg/kg (e.g., 0.1-20mg/kg) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays. An exemplary dosing regimen is disclosed in WO 94/04188.

The antibody composition will be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The "therapeutically effective amount" of the antibody to be administered will be governed by such considerations, and is the minimum amount necessary to prevent, ameliorate, or treat the LFA-1-mediated disorder, including treating rheumatoid arthritis, reducing inflammatory responses, inducing tolerance of immunostimulants, preventing an immune response that would result in rejection of a graft by a host or vice-versa, or prolonging survival of a transplanted graft. Such amount is preferably below the amount that is toxic to the host or renders the host significantly more susceptible to infections.

The antibody need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. For example, in rheumatoid arthritis, the antibody may be given in conjunction with a glucocorticosteroid. In addition, T cell receptor peptide therapy is suitably an adjunct therapy to prevent clinical signs of autoimmune encephalomyelitis. For transplants, the antibody may be administered concurrently with or separate from an immunosuppressive agent as defined above, e.g., cyclosporin A, to modulate the immunosuppressant effect. Alternatively, or in addition, VLA-4 antagonists or other LFA-1 antagonists may be administered to the mammal suffering from a LFA-1 mediated disorder. The effective amount of such other agents depends on the amount of anti-CD11a antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the heretofore employed dosages.

G. Articles of Manufacture

In another embodiment of the invention, an article of manufacture containing materials useful for the treatment of the disorders described above is provided. The article of manufacture comprises a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is

effective for treating the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agent in the composition is the anti-CD11a antibody. The label on, or associated with, the container indicates that the composition is used for treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

EXAMPLE

PRODUCTION OF HUMANIZED ANTI-CD11a ANTIBODIES

This example describes the humanization and *in vitro* biological efficacy of a murine anti-human CD11a monoclonal antibody, MHM24 (Hildreth *et al.*, *Eur. J. Immunol.* 13:202-208 (1983)). Previous studies on the murine MHM24 have shown that it, like other anti-CD11a antibodies, can inhibit T cell function (Hildreth *et al.*, *J. Immunol.* 134:3272-3280 (1985); Dougherty *et al.*, *Eur. J. Immunol.* 17:943-947 (1987)). Both the murine and humanized MAbs effectively prevent adhesion of human T cells to human keratinocytes and the proliferation of T cells in response to nonautologous leukocytes in the mixed lymphocytes response (MLR), a model for responsiveness to MHC class II antigens (McCabe *et al.*, *Cellular Immunol.* 150:364-375 (1993)). However, both the murine (Reimann *et al.*, *Cytometry*, 17:102-108 (1994)) and humanized MAbs did not cross-react with nonhuman primate CD11a other than chimpanzee CD11a. In order to have a humanized MAb available for preclinical studies in rhesus, the humanized MAb was re-engineered to bind to rhesus CD11a by changing four residues in one of the complementarity-determining regions, CDR-H2, in the variable heavy domain. Cloning and molecular modeling of the rhesus CD11a I-domain suggested that a change from a lysine residue in human CD11a I-domain to glutamic acid in rhesus CD11a I-domain is the reason that the murine and humanized MAbs do not bind rhesus CD11a.

Materials and Methods

(a) Construction of humanized F(ab')s

The murine anti-human CD11a MAb, MHM24 (Hildreth *et al.*, *Eur. J. Immunol.* 13:202-208 (1983); Hildreth *et al.*, *J. Immunol.* 134:3272-3280 (1985)), was cloned and sequenced. In order to have a plasmid useful for mutagenesis as well as for expression of F(ab)s in *E. coli*, the phagemid pEMX1 was constructed. Based on the phagemid pb0720, a derivative of pB0475 (Cunningham *et al.*, *Science* 243:1330-1336 (1989)), pEMX1 contains a DNA fragment encoding a humanized κ -subgroup I light chain and a humanized subgroup III heavy chain (VH-CH1) and an alkaline phosphatase promotor and Shine-Dalgarno sequence both derived from another previously described pUC119-based plasmid, pAK2 (Carter *et al.*, *Proc. Natl. Acad. Sci. USA* 89:4285 (1992)). A unique SpeI restriction site was also introduced between the DNA encoding for the F(ab) light and heavy chains.

To construct the first F(ab) variant of humanized MHM24, F(ab)-1, site-directed mutagenesis (Kunkel, *Proc. Natl. Acad. Sci. USA* 82:488 (1985)) was performed on a deoxyuridine-containing template of pEMX1; all six CDRs were changed to the MHM24 sequence. All other F(ab) variants were constructed from a template of F(ab)-1. Plasmids were transformed into *E. coli* strain XL-1 Blue (Stratagene, San Diego, CA) for preparation of double- and single-stranded DNA. For each variant both light and heavy chains were completely

sequenced using the dideoxynucleotide method (Sequenase, U.S. Biochemical Corp.). Plasmids were transformed into *E. coli* strain 16C9, a derivative of MM294, plated onto LB plates containing 5 µg/ml carbenicillin, and a single colony selected for protein expression. The single colony was grown in 5 ml LB-100 µg/ml carbenicillin for 5-8 h at 37°C. The 5 ml culture was added to 500 ml AP5-100 µg/ml carbenicillin and allowed to grow for 16 h in a 4 L baffled shake flask at 37°C. APS media consists of: 1.5 g glucose, 11.0 Hycase SF, 0.6 g yeast extract (certified), 0.19 g MgSO₄ (anhydrous), 1.07 g NH₄Cl, 3.73 g KCl, 1.2 g NaCl, 120 ml 1 M triethanolamine, pH 7.4, to 1 L water and then sterile filtered through 0.1 µm Sealkeen filter.

Cells were harvested by centrifugation in a 1 L centrifuge bottle (Nalgene) at 3000 x g and the supernatant removed. After freezing for 1 h, the pellet was resuspended in 25 ml cold 10 mM MES-10 mM EDTA, pH 5.0 (buffer A). 250 µl of 0.1 M PMSF (Sigma) was added to inhibit proteolysis and 3.5 ml of stock 10 mg/ml hen egg white lysozyme (Sigma) was added to aid lysis of the bacterial cell wall. After gentle shaking on ice for 1 h, the sample was centrifuged at 40,000 x g for 15 min. The supernatant was brought to 50 ml with buffer A and loaded onto a 2 ml DEAE column equilibrated with buffer A. The flow-through was then applied to a protein G-Sepharose CL-4B (Pharmacia) column (0.5 ml bed volume) equilibrated with buffer A. The column was washed with 10 ml buffer A and eluted with 3 ml 0.3 M glycine, pH 3.0, into 1.25 ml 1 M Tris, pH 8.0. The F(ab) was then buffer exchanged into PBS using a Centricon-30 (Amicon) and concentrated to a final volume of 0.5 ml. SDS-PAGE gels of all F(ab)s were run to ascertain purity and the molecular weight of each variant was verified by electrospray mass spectrometry.

(b) Construction of chimeric and humanized IgG

For generation of human IgG1 variants of chimeric (chIgG1) and humanized (HuIgG1) MHM24, the appropriate murine or humanized variable light and variable heavy (F(ab)-8, Table II) domains were subcloned into separate previously described pRK vectors (Gorman *et al.*, *DNA Protein Eng. Tech.* 2:3 (1990)). Alanine-scan variants were constructed by site-directed mutagenesis (Kunkel, *Proc. Natl. Acad. Sci. USA* 82:488 (1985)) of the HuIgG1 light and heavy chain plasmids. The DNA sequence of each variant was verified by dideoxynucleotide sequencing.

Heavy and light chain plasmids were cotransfected into an adenovirus-transformed human embryonic kidney cell line, 293 (Graham *et al.*, *J. Gen. Virol.* 36:59 (1977)), using a high efficiency procedure (Graham *et al.*, *J. Gen. Virol.* 36:59 (1977); Gorman *et al.*, *Science*, 221:551 (1983)). Media was changed to serum-free and harvested daily for up to 5 days. Antibodies were purified from the pooled supernatants using protein A-Sepharose CL-4B (Pharmacia). The eluted antibody was buffer exchanged into PBS using a Centricon-30 (Amicon), concentrated to 0.5 ml, sterile filtered using a Millex-GV (Millipore) and stored at 4°C.

The concentration of antibody was determined using total Ig-binding ELISA. The concentration of the reference humanized anti-p185^{HER}IgG1 (Carter *et al.*, *Proc. Natl. Acad. Sci. USA* 89:4285 (1992)) was determined by amino acid composition analysis. Each well of a 96-well plate was coated with 1 µg/ml of goat anti-human IgG F(ab')₂ (Cappel Laboratories, Westchester, PA) for 24 h at 4°C. Purified antibodies were diluted and added in duplicate to the coated plates. After 1.5 h incubation, the plates were washed with PBS-0.02% Tween 20 and 0.1 ml of a 1:2000 dilution of horseradish peroxidase-conjugated sheep anti-human IgG F(ab')₂ (Cappel) was added. After 1.5 h incubation the plates were washed and 0.1 ml 0.2 mg/ml o-

phenylenediamine dihydrochloride-0.01% hydrogen peroxide-PBS was added. After 10 min. the reaction was stopped with 2 M sulfuric acid and the O.D. 490 nm was read.

(c) *Cloning of rhesus CD11a I-domain*

The DNA sequence of the rhesus I-domain was obtained using RT-PCR and primers derived from the human CD11a DNA sequence. Briefly, mRNA was isolated from $\sim 10^7$ rhesus leukocytes using the Fast Track mRNA purification kit (Invitrogen). 10 μ g mRNA was reverse transcribed using MuLV reverse transcriptase. The first strand cDNA was then amplified by 40 cycles of PCR using the primers:
5'-CACTTTGGATACCGCGTCCTGCAGGT-3' (forward) (SEQ ID NO:21) and
5'-CATCCTGCAGGTCTGCCTTCAGGTCA-3' (reverse) (SEQ ID NO:22).

A single band of the predicted size was purified from the PCR reaction using agarose gel electrophoresis. The PCR product was digested with restriction enzyme Sse8387I (Takara) and ligated to a human CD11a-containing plasmid digested with the same restriction enzyme. There are two Sse8387I sites in the human CD11a sequence, one on either side of the I-domain. The resulting plasmid encoded a chimera consisting of human CD11a with a rhesus I-domain substituted for the human I-domain. DNA sequence analysis revealed five amino acid differences between human and rhesus. One difference was in the region N-terminal to the I-domain (Thr59Ser) and the other four differences were in the I-domain itself: Val133Ile, Arg189Gln, Lys197Glu, and Val308Ala (Fig. 2).

(d) *FACScan analysis of F(ab) and IgG binding to Jurkat cells*

Aliquots of 10^6 Jurkat T-cells were incubated with serial dilutions of humanized and control antibodies in PBS-0.1% BSA-10 mM sodium azide for 45 min at 4°C. The cells were washed and then incubated in fluorescein-conjugated goat anti-human F(ab')₂ (Organon Teknika, Westchester, PA) for 45 min at 4°C. Cells were washed and analysed on a FACScan (Becton Dickinson, Mountain View, CA). 8×10^3 cells were acquired by list mode and gated by forward light scatter versus side light scatter, thereby excluding dead cells and debris.

(e) *Saturation binding to determine apparent K_d s*

Radiolabeled antibodies were prepared using Iodo-Gen (Pierce, Rockford, IL) according to the manufacturer's instructions. 50 μ g of antibody and 1 mCi ¹²⁵I (DuPont, Wilmington, DE) were added to each tube and incubated for 15 min at 25°C. Radiolabeled proteins were purified from the remaining free ¹²⁵I using PD-10 columns (Pharmacia, Uppsala, Sweden) equilibrated in Hank's Balanced Salt Solution (HBSS, Life Technologies, Grand Island, NY) containing 0.2% gelatin.

Mononuclear cells were purified from heparinized human peripheral blood collected from two donors using Lymphocyte Separation Medium (LSM, Organon Teknika, Durham, NC) according to the manufacturer's instructions. The blood was centrifuged at 400 x g for 40 min at 25°C with no braking. Cells at the interface of the LSM and plasma were harvested and then resuspended in HBSS-0.2% gelatin.

Leukocytes were purified from heparinized rhesus monkey peripheral blood collected from two individuals by Dextran sedimentation. Blood was diluted with an equal volume of 3% Dextran T500 (Pharmacia) in PBS and was allowed to sediment undisturbed at 25°C for 30 min. After sedimentation, cells remaining in suspension were harvested and pelleted by centrifugation at 400 x g for 5 min. Residual

erythrocytes were removed by two cycles of hypotonic lysis using distilled water and 2X HBSS. After erythrocyte lysis, cells were washed in PBS and then resuspended in HBSS-0.2% gelatin.

Antibody affinities were determined by saturation binding using either peripheral blood mononuclear cells (murine MHM24 and HuIgG1) or rhesus leukocytes (MHM23, RhIgG1). In each assay, a radiolabeled antibody was serially diluted in HBSS-0.2% gelatin in quadruplicate. Nonspecific binding was determined by the addition of 500nM final concentration of homologous unlabeled antibody in duplicate through the serial dilution. Human lymphocytes or rhesus leukocytes were added to the plates in a volume of 170 μ l per well. Plates were incubated for 2 hr at room temperature on an orbital plate shaker. After incubation, cells were harvested using a SKATRON™ cell harvester (Lier, Norway) and washed 10 times with PBS containing 0.25% gelatin and 0.1% sodium azide. Samples were then counted for 1 min in an LBK Wallac GammaMaster gamma counter (Gaithersburg, MD). Data was transformed from counts per minute to nanomolarity and four-parameter curve-fitting of saturation plots (bound versus total) was then performed to determine K_d (app) values.

(f) *Keratinocyte monolayer adhesion assay*

Normal human epidermal keratinocytes (Clonetics, San Diego, CA) were removed from culture flasks with trypsin-EDTA, centrifuged, and resuspended in lymphocyte assay medium (RPMI 1640 (GIBCO)-10% fetal calf serum-1% penicillin/streptomycin) at a concentration of 5×10^5 viable cells/ml. Aliquots of 0.1 ml/well were then cultured overnight in flat-bottom 96-well plates; appropriate wells were stimulated by addition of interferon-gamma (Genentech, South San Francisco, CA) at 100 units/well.

Jurkat clone E6-1 cells (ATCC, Rockville, MD) or purified rhesus lymphocytes (see MLR methods) were labeled with 20 μ g/ml Calcein AM (Molecular Probes, Eugene, OR) at 37°C for 45 min. After washing three times with lymphocyte assay medium, Jurkat or rhesus lymphocyte cells were resuspended to 1×10^6 cells/ml and incubated with serially-diluted antibody at 4°C for 30 min. After removal of medium from the keratinocyte monolayer, 0.1 ml/well of labeled cells were added and incubated at 37°C for 1 h. The wells were washed five times with 0.2 ml/well/wash of 37°C lymphocyte medium to remove non-attached cells. Fluorescence was measured using a Cytofluor 2300 (Millipore, Bedford, MA).

A rhesus-human chimeric CD11a (Rh/HuCD11a) comprising human CD11a with a rhesus I-domain was constructed by site-directed mutagenesis (Kunkel, *Proc. Natl. Acad. Sci. USA* 82:488 (1985)) on a deoxyuridine-containing template plasmid encoding human CD11a. Four residues were altered: Val133Ile, Arg189Gln, Lys197Glu, and Val308Ala (Fig. 2). Plasmids coding for Rh/HuCD11a and human CD11b (EP 364, 690) were cotransfected into an adenovirus-transformed human embryonic kidney cell line, 293 (Graham *et al.*, *J. Gen. Virol.* 36:59 (1977)), using a high efficiency procedure (Graham *et al.*, *J. Gen. Virol.* 36:59 (1977); Gorman *et al.*, *Science*, 221:551 (1983)). Rh/HuCD11a-transfected 293 cells were labeled with 20 μ g/ml Calcein AM at 37°C for 45 min. After washing three times with lymphocyte assay medium, Rh/HuCD11a-transfected 293 cells were resuspended to 1×10^6 cells/ml and incubated with serially-diluted antibody at 4°C for 30 min. After removal of medium from the keratinocyte monolayer, 0.1 ml/well of labeled 293 cells was added and incubated at 37°C for 1 h. The wells were washed five times with 0.2 ml/well/wash of 37°C lymphocyte medium to remove non-attached cells. Fluorescence was measured using a Cytofluor 2300.

(g) *ICAM adhesion assay*

Maxisorp (Nunc) 96-well plates were coated with 0.1 ml/well of 1 µg/ml goat anti-human IgG Fc (Caltag) for 1 h at 37°C. After washing the plates three times with PBS, the plates were blocked with 1% BSA-PBS for 1 h at 25°C. The plates were then washed three times with PBS and 0.1 ml/well of 50 ng/ml recombinant human ICAM-IgG was added and incubated overnight.

The ICAM-IgG consisted of the five extracellular domains of human ICAM fused onto a human IgG Fc. A plasmid for the expression of a human ICAM-1 (Simmons *et al. Cell* 331:624-627 (1988) and Staunton *et al. Cell* 52:925-933 (1988)) immunoadhesin called pRK.5dICAMGalg was constructed. It contains the five Ig-like domains of ICAM-1, a six amino acid cleavage site recognized by an H64A variant of subtilisin BPN', Genenase I (Carter *et al. Proteins: Structure, Function and Genetics* 6:240-248 (1989)), and the Fc region from human IgG1 (Ellison *et al. Nucleic Acids Research* 10:4071-4079 (1982)) in the pRK5 vector (Eaton *et al. Biochemistry* 25:8343-8347 (1986)). Human embryonic kidney 293 cells (Graham *et al. J. Gen. Virol.* 36:59 (1977)) were stably transfected with pRK.5dICAMGalg and the RSV-neo plasmid (Gorman *et al. Science* 221:551-553 (1983)) to generate a cell line expressing the five domain ICAM Ig (5dICAMIg). A clone was selected which expressed 20 µg/ml of secreted 5dICAMIg by enzyme-linked immunosorbent assay (ELISA), using antibodies to human IgG Fc (Caltag, Burlingame, CA) and ICAM-1 (BBIG-I1; R & D Systems, Minneapolis, MN). Cell culture supernatant from this cell line was loaded onto a Protein A column (ProsepA, Bioprocessing, Ltd., Durham, England) equilibrated in 0.01 M Hepes buffer (pH 7.0), 0.15 M NaCl (HBS) and the column was washed with HBS followed by 0.01 M Hepes buffer (pH 7.0), 0.5 M NaCl, 0.5 M TMAC (tetra-methyl ammonium chloride) to remove non-specifically bound material. The TMAC buffer was thoroughly washed from the column with HBS and the 5dICAMIg eluted with 0.01 M Hepes buffer (pH 7.0), 3.5 M MgCl₂ and 10% (w/v) glycerol. The protein A pool was dialyzed extensively against HBS and concentrated.

Purified rhesus lymphocytes (see MLR methods) were labeled with 20 µg/ml Calcein AM (Molecular Probes, Eugene, OR) at 37°C for 45 min. After washing three times with lymphocyte assay medium, rhesus lymphocyte cells were resuspended to 1×10^6 cells/ml and incubated with serially-diluted antibody at 4°C for 30 min. After removal of medium from the ICAM-IgG coated plates, 0.1 ml/well of labeled cells were added and incubated at 37°C for 1 h. The wells were washed five times with 0.2 ml/well/wash of 37°C lymphocyte medium to remove non-attached cells. Fluorescence was measured using a Cytofluor 2300 (Millipore, Bedford, MA).

(h) *One-way mixed lymphocyte response (MLR)*

For both human and rhesus MLR, peripheral blood lymphocytes from two unrelated donors were isolated from whole, heparinized blood using Lymphocyte Separation Medium (Organon Teknika, Durham, NC). Lymphocytes were resuspended to a concentration of 3×10^6 cells/ml in RPMI 1640 (GIBCO)-10% human AB serum-1% glutamine-1% penicillin/streptomycin-1% non-essential amino acids-1% pyruvate- 5×10^{-5} M 2-β-mercaptoethanol-50 µg/ml gentamycin-5 µg/ml polymyxin B. The stimulator cells were made unresponsive by irradiation with 3000 rads in a cesium irradiator. Responder cells at a concentration of 1.5×10^5 cells per well were co-cultured with an equal number of stimulator cells in 96-well, flat-bottom plates. Serial two-fold dilutions of each antibody were added to the cultures to give a total volume of 200 µl/well. The

cultures were incubated at 37°C in 5% CO₂ for 5 days and then pulsed with 1 µCi/well of [³H]thymidine for 16 h. [³H]thymidine incorporation was measured with a Beckman scintillation counter. Assays were done in triplicate. A humanized anti-human p185^{HER2} MAb (Carter *et al.*, *Proc. Natl. Acad. Sci. USA* 89:4285 (1992)) was used as isotype control for HuIgG1 and RhIgG1. A murine anti-hamster tPA MAb (Genentech) was used as isotype control (murine IgG1) for the MHM23 MAb. MAb 25.3 was purchased from Immunotech, Inc. (Westbrook, ME).

(i) *Computer graphics models of murine and humanized MHM24*

Sequences of the VL and VH domains (Figs. 1A & B) were used to construct a computer graphics model of the murine MHM24 VL-VH domains. This model was used to determine which framework residues should be incorporated into the humanized antibody. A model of F(ab)-8 was also constructed to verify correct selection of murine framework residues. Construction of the models was performed as described previously (Carter *et al.*, *Proc. Natl. Acad. Sci. USA* 89:4285 (1992); Eigenbrot *et al.*, *J. Mol. Biol.* 229:969 (1993)).

Results

(a) *Humanization*

The consensus sequence for the human heavy chain subgroup III and the light chain subgroup κ I were used as the framework for the humanization (Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) (Figs. 1A & B). This framework has been successfully used in the humanization of other murine antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci. USA* 89:4285 (1992); Presta *et al.*, *J. Immunol.* 151:2623-2632 (1993); Eigenbrot *et al.*, *Proteins* 18:49-62 (1994)). All humanized variants were initially made and screened for binding as F(ab)s expressed in *E. coli*. Typical yields from 500 ml shake flasks were 0.2-0.5 mg F(ab). Mass spectrometry verified the mass of each F(ab) to within 5 mass units.

CDR-H1 included residues H28-H35, which includes all exposed residues from both Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991) and Chothia *et al.* *Nature* 342:877-883 (1989). The other hypervariable loops were defined according to Chothia *et al.* (1989). Light chain residue numbers are prefixed with L; heavy chain residue numbers are prefixed with H.

Table II

Binding of humanized MHM24 variants to human CD11a on Jurkat cells

	Variant	Template	Changes ^a	Purpose	EC50 F(ab)/ EC50 F(ab)-2 ^b		
					Mean	S.D.	N
	F(ab)-1	Human FR	ArgH71Val	Straight CDR swap	no binding		2
5	F(ab)-2	F(ab)-1	AlaH60Asn AspH61Gln SerH62Lys ValH63Phe GlyH65Asp	Extended CDR-H2 (Kabat <i>et al.</i> (1991))	1.0		4
	F(ab)-3	F(ab)-2	PheH67Ala	Packing;CDR-H2	1.2	0.33	3
	F(ab)-4	F(ab)-2	ValH71Arg	Packing;CDR-H1,H2	2.9		1
10	F(ab)-5	F(ab)-2	AsnH73Lys LysH75Ser AsnH76Ser	Framework loop in VH	0.043	0.015	4
	F(ab)-6	F(ab)-5	SerL53Thr GluL55Gln	Extended CDR-L2	0.012	0.005	4
	F(ab)-7	F(ab)-6	PheH27Tyr	Extended CDR-H1	0.004	0.002	4
15	F(ab)-8	F(ab)-7	SerH75Lys SerH76Asn	Framework loop in VH back to human	0.004	0.002	4
	HuIgG1 ^c				0.004	0.002	4
	chIgG1 ^d				0.006	0.005	4

^a Murine residues are in bold; residue numbers are according to Kabat *et al.*, *Sequences of Proteins of Immunological Interest* 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991).

^b Mean and standard deviation are the average of the ratios calculated for each of the independent FACSscan assays; the EC50 for F(ab)-2 was 771 ± 320 ng/ml.

^c HuIgG1 is F(ab)-8 VL and VH domains fused to human constant light and heavy chains.

^d chIgG1 is chimeric IgG with murine VL and VH domains fused to human constant light and heavy chains.

In the initial variant F(ab)-1 the CDR residues were transferred from the murine antibody to the human framework. In addition, residue H71 was changed from the human Arg to murine Val since this residue has been shown previously to affect the conformations of CDR-H1 and CDR-H2 (Chothia *et al.*, *Nature*, 342:877-883 (1989); Tramontano *J. Mol. Biol.* 215:175 (1990)). This F(ab) showed no detectable binding. In F(ab)-2 CDR-H2 was extended to include the sequence-based definition (*i.e.*, including residues H60-H65). The EC50 for F(ab)-2 binding to human CD11a was 771 ± 320 ng/ml, which was 148-fold weaker than the EC50 of the chimeric IgG1 (5.2 ± 3.0 ng/ml).

In previous humanizations, it has been found that residues in a framework loop (FR-3) adjacent to CDR-H1 and CDR-H2 can affect binding (Eigenbrot *et al.*, *Proteins* 18:49-62 (1994)). In F(ab)-5 three residues in this loop were changed to their murine counterpart and this variant showed a 23-fold improvement in binding (Table II). Alteration of human residues at positions L53 and L55 to murine (*i.e.* SerL53Thr and GluL55Gln) further improved binding by another 4-fold (F(ab)-6, Table II); this effectively converted CDR-L2 from the structure-based definition (residues L50-L52) to the sequence-based definition (residues L50-L56). Subsequent alteration of PheH27 to murine Tyr in CDR-H1 resulted in an additional 3-fold improvement (F(ab)-7; Table II). Finally, based on the models of murine and humanized MHM24, two of the three murine residues (H75 and H76) in FR-3 were changed back to human and it was found that these two residues had no effect on binding (cf. F(ab)-7 and F(ab)-8, Table II). The average EC50 for F(ab)-8 was slightly better than that of the chimeric IgG1 (Table II). Not all changes from human to murine resulted in improved binding. PheH67 was changed to murine Ala since this position had been previously found to affect binding (Presta *et al.*, *J. Immunol.* 151:2623-2632 (1993)) but no effect was evident (F(ab)-3, Table II). Changing ValH71 back to the human Arg effected a 3-fold reduction in binding (F(ab)-4, Table II), supporting inclusion of ValH71 in F(ab)-1.

The VL and VH domains from F(ab)-8 were transferred to human IgG1 constant domains. The full length intact antibody, HuIgG1, showed an EC50 equivalent to F(ab)-8 and improved compared to the full length chimeric IgG1 (Table II). When data for all assays of HuIgG1 is considered, including its use as a standard for the alanine-scan and MLR assays (see below), the EC50 for HuIgG1 against human CD11a was 0.042 ± 0.072 nM (N = 15). Saturation binding analysis was also performed to determine the apparent dissociation constants, $K_d(\text{app})$: 0.15 ± 0.02 nM for murine MHM24 and 0.15 ± 0.04 nM for HuIgG1 (Table III).

Table III

Apparent K_d by saturation binding to human lymphocytes and rhesus leukocytes

	muMHM24 $K_d(\text{app})$ nM	HuIgG1 $K_d(\text{app})$ nM	muMHM23 $K_d(\text{app})$ nM	RhIgG1 $K_d(\text{app})$ nM
human donor 1	0.16 +/- 0.01	0.11 +/- 0.08		
human donor 2	0.13 +/- 0.02	0.18 +/- 0.03		
rhesus donor 1			3.9 +/- 0.31	3.9 +/- 1.04
rhesus donor 2			4.5 +/- 0.51	n.d.
rhesus donor 3				2.8 +/- 1.1 ^a
rhesus donor 3				2.7 +/- 0.9

^a Assays of rhesus donor 3 were performed using two batches of RhIgG1; the assays were performed in the presence of 1 mg/ml human IgG1 to block Fc receptor interaction.

(b) *Alanine-Scan of CDR Residues*

To determine which CDR residues were involved in binding to human CD11a an alanine-scan (Cunningham *et al.*, *Science* 244:1081 (1989)) was performed on the CDR residues of HuIgG1. Each variant was tested for binding to CD11a on Jurkat cells. In the light chain only CDR-L3 contributes to binding. HisL91 had a large effect (Table IV) and is probably conformational since this side chain should be partially buried. Residues AsnL92 and TyrL94 had more modest effect, reducing binding by 3-fold and 12-fold, respectively. Note however that simultaneously changing these two residues to alanine (as well as GluL93Ala) had a non-additive effect on binding (variant L3, Table IV).

Table IV

Alanine-scan of humanized MHM24 CDR residues

10

Variant IgG1		Human CD11a Var.EC50/HuIgG1 EC50 ^b			Rhesus CD11a Var.EC50/HuIgG1 EC50 ^c		
CDR Sequence ^a		Mean	S.D	N	Mean	S.D.	N
CDR-H1	G Y S F T G H W M N						
15	H1 ^d	5.9	0.8	2	nb		
	SerH28Ala	6.9	0.1	2	nb		
	ThrH30Ala	1.7	0.3	2	1.3		
	GlyH31Ala	1.2	0.1	2	2.4		
	HisH32Ala	2.3	0.2	2	nb		
	TrpH33Ala	>870		1	nb		
20	CDR-H2						
	H2	14.1	7.8	10	0.055	0.050	15
	H2B	>600		2	nb		
	H2A1	10.8	7.3	6	0.013	0.012	10
	H2A2	1.9	0.1	2	1.1	0.1	2
25	H2A3	4.6	0.2	2	1.3		
	HisH52Ala	1.5	0.2	2	0.7		
	HisH52Ser	5.0	0.3	2	nb		
	SerH53Ala	1.8	0.1	2	0.7		
	AspH54Ala	147	18	2	0.3		
30	SerH55Ala	1.3	0.1	2	1.3		
	SerH55Asn	2.1	0.2	2	nb		
	SerH55Gln	2.9	1.4	2	6.7		
	GluH56Ala	4.0	0.6	2	0.3		
	ArgH58Ala	1.1	0.1	2	3.3		
35	GlnH61Ala	4.1	0.1	2	5.3		
	LysH62Ala	1.8	0.2	2	4.9		
	LysH64Ala	2.5	0.1	2	1.2		
	AspH65Ala	0.8	0.1	2	1.1		
40	FR-3						
	LysH73Ala	5.2	0.9	2	nb		
	LysH73Arg	4.6	1.1	2	5.5		
45	CDR-H3						
	H3	>900		2	nb		
	H3B	34.7	13.6	2	nb		
	TyrH97Ala	10.9	2.1	2	nb		
	TyrH99Ala	1.4	0.1	2	nb		
	ThrH100aAla	2.3	0.6	2	nb		

Variant IgG1		Human CD11a Var.EC50/HuIgG1 EC50 ^b			Rhesus CD11a Var.EC50/HuIgG1 EC50 ^c		
CDR Sequence ^a		Mean	S.D	N	Mean	S.D.	N
ThrH100bAla	A	1.4	0.2	2	nb		
TyrH100cAla	A	7.6	1.0	2	nb		
CDR-L1	R A S K T I S K Y L A						
L1	AA AA	1.3	0.3	3	1.0		
CDR-L2	S G S T L Q S						
L2	A AA	1.1	0.0	2	nb		
SerL50Ala	A	1.2	0.5	2	2.7		
SerL52Ala	A	1.1	0.2	2	13.3		
ThrL53Ala	A	0.9	0.4	2	0.7		
CDR-L3	Q Q H N E Y P L T						
L3	AAA	>900		2	nb		
HisL91Ala	A	>900		2	nb		
AsnL92Ala	A	3.3	0.7	2	3.3		
GluL93Ala	A	1.7	0.2	2	2.9		
TyrL94Ala	A	11.8	0.1	2	nb		
hu4D5 ^e		>900		5	nb		

^a CDRs and FR-3 are as defined in Kabat *et al.*, (1991) *supra*.

^b EC50 HuIgG1 for human CD11a = 0.042 nM (S.D.= 0.072; N = 15).

^c EC50 HuIgG1 for rhesus CD11a = 45.6 nM (S.D.= 40.4; N = 16); all values for rhesus CD11a are for a single binding assay unless otherwise noted; nb denotes binding of variant is greater than 10-fold weaker than HuIgG1.

^d Multiple alanine variants:

H1, SerH28Ala/ThrH30Ala/HisH32Ala;
H2, HisH52Ala/SerH53Ala/SerH55Ala;
H2B, AspH54Ala/GluH56Ala/ArgH58Ala;
H2A1, HisH52Ala/SerH53Ala;
H2A2, SerH53Ala/SerH55Ala;
H2A3, HisH52Ala/SerH55Ala;
H3, TyrH97Ala/TyrH99Ala;
H3B, ThrH100aAla/ThrH100bAla/TyrH100cAla;
L1, LysL27Ala/ThrL28Ala/SerL30Ala/LysL31Ala;
L2, SerL50Ala/SerL52Ala/ThrL53Ala;
L3 AsnL92Ala/GluL93Ala/TyrL94Ala.

^e hu4D5 is a humanized anti-p185^{HER2} antibody with the same IgG1 framework as the huMHM24 antibody (Carter *et al.*, *Proc.Natl. Acad. Sci. USA* 89:4285 (1992)).

In the heavy chain, CDR-H2 and CDR-H3 are the prominent contributors to the binding. CDR-H1 residue TrpH33Ala had a large effect but this is most likely due to a conformational change as TrpH33 should be partially buried. The most important single residue contributing to the binding is AspH54 in CDR-H2; changing this residue to alanine effected a 147-fold reduction in binding (Table IV). Other residues in CDR-H2 involved in binding include GluH56, GlnH61 and LysH64 (Table IV). In CDR-H3, TyrH97Ala reduced binding by 11-fold and TyrH100cAla by 8-fold. As in CDR-L3, simultaneous alteration of several CDR-H3 residues to alanine effected a non-additive, large reduction in binding (cf. variant H3 versus TyrH97Ala and

TyrH99Ala, Table IV). In addition, the FR-3 residue included in the humanization, LysH73, also showed a 5-fold reduction in binding when changed to alanine or arginine (Table IV).

(c) *Re-engineering HuIgG1 to Bind to Rhesus CD11a*

Both murine MHM24 and HuIgG1 showed approximately 1000-fold reduction in binding to rhesus CD11a: HuIgG1 had an EC50 against rhesus CD11a of 45.6 ± 40.4 nM (N = 16) compared to an EC50 of 0.042 ± 0.072 nM against human CD11a. Since a primate model is important in evaluating the biology, toxicity, and efficacy of MHM24, improving the binding of HuIgG1 to rhesus CD11a was deemed advantageous. Initially, the MAb hypervariable region residues which were important in binding to human CD11a and rhesus CD11a were determined so that those important for the rhesus but not the human could be altered. Accordingly, the alanine-scan variants were also assayed against rhesus CD11a on peripheral blood lymphocytes. The most important finding was that one of the multiple-alanine mutation variants, variant H2, bound 18-fold better to rhesus CD11a than HuIgG1 (Table IV). However, individual mutations at the three residues included in variant H2 showed minimal improvement in binding: HisH52Ala, 0.7-fold better, SerH53Ala, 0.7-fold better, and SerH55Ala, 1.3-fold worse (Table IV). A series of double mutations at these three residues showed that the combination HisH52Ala-SerH53Ala was the best, providing a 77-fold improvement in binding compared to HuIgG1 (cf. variants H2A1, H2A2 and H2A3, Table IV). In addition, the AspH54Ala and GluH56Ala variants also effected a 3-fold improvement over HuIgG1 (Table IV), even though AspH54 is the most important binding residue in HuIgG1 with respect to human CD11a.

In an attempt to find a single substitution at position H54 which would improve binding to rhesus CD11a but not reduce binding to human CD11a, position H54 was substituted with a variety of amino acids. All substitutions reduced binding by greater than an order of magnitude whereas the substitution AspH54Asn improved rhesus binding by 10-fold (Table V).

Table V
Amino acid substitution at AspH54

AspH54 change to	Variant EC50/HuIgG1 EC50 ^a		
	Human CD11a ^b		Rhesus CD11a ^c
	mean	S.D.	
Ala	147	18	0.3
Asn	26	1	0.1
Gln	20	1	4.4
Glu	16	2	>25
Ser	>100		0.9
Arg	>250		>25
Lys	>100		3.8
His	>300		>25
Thr	>450		>25

Met	>150		>25
Leu	>300		>25

^a EC50 HuIgG1 for human CD11a = 0.042 nM (S.D.= 0.072; N=15); EC50 HuIgG1 for rhesus CD11a = 45.6 nM (S.D.= 40.4; N=16).

5 ^b Values are the mean of two assays.

^c Values are for a single assay.

Since non-additive effects were noted for changes at positions H52-H53, these were combined with a variety of changes at positions H54 and H56 (Table VI). For all variants, H52 and H53 were alanine. In one series, position H54 was Asn and position H56 was Glu (original), Ala, Asn or Gln. None of these variants improved rhesus CD11a binding over the H2A1 variant (Table VI). In another series, position H54 was Ala and position H56 was Glu (original), Ala, Ser or Asn and again all were worse than variant H2A1. In the third series, position H54 was Ser and position H56 was Glu (original), Ala, Ser or Asn. Two of these variants exhibited improved binding to rhesus CD11a compared to the H2A1 variant (H2C11 and H2C12, Table VI). The rhesus CD11a EC50 for these two variants was 0.11 ± 0.11 nM (N = 9) for H2C11 and 0.19 ± 0.08 nM (N = 7) for H2C12. These values are 3- to 5-fold weaker than the EC50 of HuIgG1 for human CD11a (0.042 nM) but are a 240- to 415-fold improvement over the EC50 of HuIgG1 for rhesus CD11a (45.6 nM). H2C12 will hereafter be referred to as RhIgG1. Apparent K_d values from saturation binding experiments showed that RhIgG1 bound to rhesus CD11a as well as murine MHM23 bound to rhesus CD18 (Table III).

Table VI

Binding of CDR-H2 variants to human and rhesus CD11a

20

25

30

Variant IgG1	Sequence	Var.EC50/HuIgG1 EC50 ^a		
		Human CD11a	Rhesus CD11a	
			Mean	S.D.
H2C2	A	2.6	>100	0.012(N=10)
H2C3	A A A N	>100	>100	
CDR-H2	M I H P S D S E T R Y	1.0	1.00	
H2A1	A A	10.8	0.013	
H2C1	A A N	>100	0.56	
H2C4	A A N A	>100	0.38	
H2C5	A A N N	46	0.11	
H2C6	A A N Q	>100	0.21	
H2C8	A A A	12.7	0.38	
H2C7	A A A A	2.4	1.03	
H2C10	A A A S	14.2	0.22	0.03

5	H2C9	A	A	A	N	34.3	0.22	0.04
	H2C13	A	A	S			0.10	0.06
	H2C14	A	A	S	A		0.021	0.013
	H2C12	A	A	S	S		0.004	0.001 (N=7)
	H2C11	A	A	S	N	24.9	0.002	0.001(N=9)

^a EC50 HuIgG1 for human CD11a = 0.042 nM (S.D.= 0.072; N = 15); EC50 HuIgG1 for rhesus CD11a = 45.6 nM (S.D.= 40.4; N = 16); all values for rhesus CD11a are the mean of two independent binding assays except as noted.

For the HuIgG1-human CD11a interaction, AspH54 was the most important residue (Table IV); changing this residue to other amino acids significantly reduced binding with the least reduction occurring for changes to Glu, Asn and Gln. However, for the HuIgG1-rhesus CD11a interaction, AspH54 was deleterious since changing this residue to Ala or Asn improved binding (Table V). In order to understand this difference between binding to human and rhesus CD11a, the latter was cloned from a rhesus PBL library. Fig. 2 shows that rhesus CD11a I-domain differs from human CD11a I-domain at only four positions: 133, 189, 197, 308. Previously the human CD11a epitope of MHM24 was mapped to residues 197-203 (Champe *et al.*, *J. Biol. Chem.* 270:1388-1394 (1995)) which includes the human Lys197 to rhesus Glu197 change in rhesus.

(d) *Keratinocyte Cell Adhesion Assays*

Murine MHM24, chimeric IgG1 and HuIgG1 were compared in their ability to prevent adhesion of Jurkat cells (human T-cells expressing LFA-1) to normal human epidermal keratinocytes expressing ICAM-1. All three antibodies performed similarly (Fig. 3) with similar IC50 values (Table VII).

Table VII

Blocking of cell adhesion by MHM24 variants

mAb	IC50 Value (nM)			
	Jurkat:HuK ^a	RhLy ^b :HuK Mean S.D. N	RhLy:HuICAM ^c Mean S.D. N	Rh/HuCD11a ^d :HuK Mean S.D. N
murMHM24	0.09			
HuIgG1	0.13			
chIgG1	0.10			
RhIgG1		119 86 4	5.3 4.5 3	4.9 0.2 2
MHM23		1.6 1.5 3	1.2	1.4 0.1 2

^a HuK = normal human epidermal keratinocyte.

^b RhLy = rhesus lymphocyte.

^c HuICAM = recombinant human ICAM-1.

^d Rh/HuCD11a = human CD11a with rhesus I-domain transfected into human 293 cells.

Neither murine nor humanized MHM24 blocked rhesus or cynomolgus lymphocytes from adhering to human keratinocytes. When RhIgG1 was compared to the murine anti-human CD18 antibody MHM23

(Hildreth *et al.*, *Eur. J. Immunol.* 13:202-208 (1983); Hildreth *et al.*, *J. Immunol.* 134:3272-3280 (1985)) in blocking adherence of rhesus lymphocytes to human keratinocytes, RhIgG1 was 74-fold less efficacious than MHM23 (Fig. 4A, Table VII). However, when recombinant human ICAM-1 was coated on plates (instead of human keratinocytes) RhIgG1 was only 4-fold less efficacious than MHM23 (Fig. 4B, Table VII). A chimeric CD11a comprised of human CD11a in which the I-domain was mutated to rhesus (Val133Ile, Arg189Gln, Lys197Glu, Val308Ala) was transfected into human embryonic kidney 293 cells. Again, RhIgG1 was only 4-fold down from MHM23 in blocking these Rh/HuCD11a-293 cells from adhering to human keratinocytes (Fig. 4C, Table VII).

Control isotype antibodies for RhIgG1 (humanized anti-p185HER2 antibody; Carter *et al.*, *Proc. Natl. Acad. Sci. USA* 89:4285 (1992)) and MHM23 (murine MAb 354, a murine IgG1 anti-hamster tPA) did not block binding of rhesus lymphocytes to human keratinocytes or recombinant ICAM-1 (Figs. 4A, 4B) or Rh/HuCD11a to human keratinocytes (Fig. 4C). This implies that the reduced performance of RhIgG1 compared to murine MHM23 in the rhesus lymphocyte:human keratinocyte assay was not due to any unexpected interaction of the human Fc of HuIgG1 (compared to the murine Fc of MHM23) with the rhesus lymphocytes, which might reduce the concentration of RhIgG1 available for binding to rhesus CD11a. The recombinant human ICAM-1 data show that RhIgG1 is binding to the rhesus lymphocytes and preventing adherence almost as well as murine MHM23 (Fig. 4B, Table VII). The Rh/HuCD11a-293 data (Fig. 4C, Table VII) show that RhIgG1 is not binding to targets on the human keratinocytes (compared to HuIgG1), which might reduce the concentration of RhIgG1 available for binding to rhesus CD11a. In addition, the K_d (app) of RhIgG1 to rhesus leukocytes was similar with (rhesus donor 3) and without (rhesus donor 1) addition of 1 mg/ml human IgG1 (Table III); This shows that binding of RhIgG1 is specific to rhesus CD11a.

(e) *Mixed Lymphocyte Response Assays*

In the MLR, HuIgG1 exhibited an IC50 value 2-fold weaker than the murine MHM24 (Table VIII, Fig. 5).

Table VIII
Mixed lymphocyte response assay results

mAb ^a	IC50 Value (nM)		
	Mean	S.D.	N
murMHM24	0.19	0.06	3
HuIgG1	0.38	0.14	4
mAb 25.3	3.8	1.0	2
RhIgG1	23.4	11.4	2
MHM23	30.4	24.0	3

^a murMHM24, HuIgG1 and mAb 25.3 tested in human MLR; RhIgG1 and MHM23 tested in rhesus MLR.

Both the murine and humanized MABs fared 10- to 20-fold better than MAb 25.3, which has been previously tested *in vivo* (Fischer *et al.*, *Blood* 77:249-256 (1991); Stoppa *et al.*, *Transplant Intl.* 4:3-7 (1991); Hourmant *et al.*, *Transplantation* 58:377-380 (1994)). The rhesus-binding variant RhIgG1 exhibited an IC50 value slightly better than murine MHM23 (Table VIII). Different responder:stimulator blood donors were used

in independent assays and the results did not vary significantly. The K_d of RhIgG1 for rhesus CD11a is about 26-fold down from the K_d of HuIgG1 for human CD11a (Table III) and this is reflected in the IC50-values derived from the MLR assays (Table VIII).

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Genentech, Inc.

(ii) TITLE OF INVENTION: Humanized Anti-CD11a Antibodies

5 (iii) NUMBER OF SEQUENCES: 24

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Genentech, Inc.

(B) STREET: 1 DNA Way

(C) CITY: South San Francisco

10 (D) STATE: California

(E) COUNTRY: USA

(F) ZIP: 94080

(v) COMPUTER READABLE FORM:

15 (A) MEDIUM TYPE: 3.5 inch, 1.44 Mb floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: WinPatin (Genentech)

(vi) CURRENT APPLICATION DATA:

20 (A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Lee, Wendy M.

(B) REGISTRATION NUMBER: 40,378

25 (C) REFERENCE/DOCKET NUMBER: P1014PCT

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 650/225-1994

(B) TELEFAX: 650/952-9881

(2) INFORMATION FOR SEQ ID NO:1:

30 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 108 amino acids

(B) TYPE: Amino Acid

(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

35 Asp Val Gln Ile Thr Gln Ser Pro Ser Tyr Leu Ala Ala Ser Pro
1 5 10 15Gly Glu Thr Ile Ser Ile Asn Cys Arg Ala Ser Lys Thr Ile Ser
20 25 3040 Lys Tyr Leu Ala Trp Tyr Gln Glu Lys Pro Gly Lys Thr Asn Lys
35 40 45Leu Leu Ile Tyr Ser Gly Ser Thr Leu Gln Ser Gly Ile Pro Ser
50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
65 70 75

Ser Ser Leu Glu Pro Glu Asp Phe Ala Met Tyr Tyr Cys Gln Gln
80 85 90

5 His Asn Glu Tyr Pro Leu Thr Phe Gly Thr Gly Thr Lys Leu Glu
95 100 105

Leu Lys Arg
108

(2) INFORMATION FOR SEQ ID NO:2:

10 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

15 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
1 5 10 15

Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Lys Thr Ile Ser
20 25 30

Lys Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
20 35 40 45

Leu Leu Ile Tyr Ser Gly Ser Thr Leu Gln Ser Gly Val Pro Ser
50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
65 70 75

25 Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
80 85 90

His Asn Glu Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu
95 100 105

Ile Lys Arg
30 108

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
(B) TYPE: Amino Acid
35 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
1 5 10 15

Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser

	20	25	30
	Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys		
	35	40	45
5	Leu Leu Ile Tyr Ala Ala Ser Ser Leu Glu Ser Gly Val Pro Ser		
	50	55	60
	Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile		
	65	70	75
	Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln		
	80	85	90
10	Tyr Asn Ser Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu		
	95	100	105
	Ile Lys Arg		
	108		

(2) INFORMATION FOR SEQ ID NO:4:

- 15 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 121 amino acids
 - (B) TYPE: Amino Acid
 - (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

20	Glu Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Met Arg Pro Gly		
	1	5	10 15
	Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr		
	20	25	30
25	Gly His Trp Met Asn Trp Val Arg Gln Arg Pro Gly Gln Gly Leu		
	35	40	45
	Glu Trp Ile Gly Met Ile His Pro Ser Asp Ser Glu Thr Arg Leu		
	50	55	60
	Asn Gln Lys Phe Lys Asp Lys Ala Thr Leu Thr Val Asp Lys Ser		
	65	70	75
30	Ser Ser Ser Ala Tyr Met Gln Leu Ser Ser Pro Thr Ser Glu Asp		
	80	85	90
	Ser Ala Val Tyr Tyr Cys Ala Arg Gly Ile Tyr Phe Tyr Gly Thr		
	95	100	105
35	Thr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val Ser		
	110	115	120
	Ser		
	121		

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 121 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1             5             10             15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Ser Phe Thr
             20             25             30

10 Gly His Trp Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
             35             40             45

Glu Trp Val Gly Met Ile His Pro Ser Asp Ser Glu Thr Arg Tyr
             50             55             60

Asn Gln Lys Phe Lys Asp Arg Phe Thr Ile Ser Val Asp Lys Ser
15             65             70             75

Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
             80             85             90

Thr Ala Val Tyr Tyr Cys Ala Arg Gly Ile Tyr Phe Tyr Gly Thr
             95             100            105

20 Thr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser
             110            115            120

Ser
121

```

(2) INFORMATION FOR SEQ ID NO:6:

25 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 113 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

30 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
    1             5             10             15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
             20             25             30

Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35             35             40             45

Glu Trp Val Ser Val Ile Ser Gly Asp Gly Gly Ser Thr Tyr Tyr
             50             55             60

Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser
             65             70             75

```

Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90

Thr Ala Val Tyr Tyr Cys Ala Arg Gly Phe Asp Tyr Trp Gly Gln
95 100 105

5 Gly Thr Leu Val Thr Val Ser Ser
110 113

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 184 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Lys Gly Asn Val Asp Leu Val Phe Leu Phe Asp Gly Ser Met Ser
1 5 10 15

15 Leu Gln Pro Asp Glu Phe Gln Lys Ile Leu Asp Phe Met Lys Asp
20 25 30

Val Met Lys Lys Leu Ser Asn Thr Ser Tyr Gln Phe Ala Ala Val
35 40 45

20 Gln Phe Ser Thr Ser Tyr Lys Thr Glu Phe Asp Phe Ser Asp Tyr
50 55 60

Val Lys Arg Lys Asp Pro Asp Ala Leu Leu Lys His Val Lys His
65 70 75

Met Leu Leu Leu Thr Asn Thr Phe Gly Ala Ile Asn Tyr Val Ala
80 85 90

25 Thr Glu Val Phe Arg Glu Glu Leu Gly Ala Arg Pro Asp Ala Thr
95 100 105

Lys Val Leu Ile Ile Ile Thr Asp Gly Glu Ala Thr Asp Ser Gly
110 115 120

30 Asn Ile Asp Ala Ala Lys Asp Ile Ile Arg Tyr Ile Ile Gly Ile
125 130 135

Gly Lys His Phe Gln Thr Lys Glu Ser Gln Glu Thr Leu His Lys
140 145 150

Phe Ala Ser Lys Pro Ala Ser Glu Phe Val Lys Ile Leu Asp Thr
155 160 165

35 Phe Glu Lys Leu Lys Asp Leu Phe Thr Glu Leu Gln Lys Lys Ile
170 175 180

Tyr Val Ile Glu
184

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 184 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

	Lys	Gly	Asn	Val	Asp	Leu	Ile	Phe	Leu	Phe	Asp	Gly	Ser	Met	Ser	
	1					5				10					15	
	Leu	Gln	Pro	Asp	Glu	Phe	Gln	Lys	Ile	Leu	Asp	Phe	Met	Lys	Asp	
					20					25					30	
10	Val	Met	Lys	Lys	Leu	Ser	Asn	Thr	Ser	Tyr	Gln	Phe	Ala	Ala	Val	
					35					40					45	
	Gln	Phe	Ser	Thr	Ser	Tyr	Lys	Thr	Glu	Phe	Asp	Phe	Ser	Asp	Tyr	
					50					55					60	
	Val	Lys	Gln	Lys	Asp	Pro	Asp	Ala	Leu	Leu	Glu	His	Val	Lys	His	
15					65					70					75	
	Met	Leu	Leu	Leu	Thr	Asn	Thr	Phe	Gly	Ala	Ile	Asn	Tyr	Val	Ala	
					80					85					90	
	Thr	Glu	Val	Phe	Arg	Glu	Glu	Leu	Gly	Ala	Arg	Pro	Asp	Ala	Thr	
					95					100					105	
20	Lys	Val	Leu	Ile	Ile	Ile	Thr	Asp	Gly	Glu	Ala	Thr	Asp	Ser	Gly	
					110					115					120	
	Asn	Ile	Asp	Ala	Ala	Lys	Asp	Ile	Ile	Arg	Tyr	Ile	Ile	Gly	Ile	
					125					130					135	
	Gly	Lys	His	Phe	Gln	Thr	Lys	Glu	Ser	Gln	Glu	Thr	Leu	His	Lys	
25					140					145					150	
	Phe	Ala	Ser	Lys	Pro	Ala	Ser	Glu	Phe	Val	Lys	Ile	Leu	Asp	Thr	
					155					160					165	
	Phe	Glu	Lys	Leu	Lys	Asp	Leu	Phe	Thr	Glu	Leu	Gln	Lys	Lys	Ile	
					170					175					180	
30	Tyr	Ala	Ile	Glu												
					184											

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Lys	His	Val	Lys	His	Met	Leu
1				5		7

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Gly Tyr Ser Phe Thr Gly His Trp Met Asn
1 5 10

(2) INFORMATION FOR SEQ ID NO:11:

- 10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

15 Met Ile His Pro Ser Asp Ser Glu Thr Arg Tyr Asn Gln Lys Phe
1 5 10 15
Lys Asp
17

(2) INFORMATION FOR SEQ ID NO:12:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

25 Gly Ile Tyr Phe Tyr Gly Thr Thr Tyr Phe Asp Tyr
1 5 10 12

(2) INFORMATION FOR SEQ ID NO:13:

- 30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Arg Ala Ser Lys Thr Ile Ser Lys Tyr Leu Ala
1 5 10 11

35 (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Ser Gly Ser Thr Leu Gln Ser
1 5 7

(2) INFORMATION FOR SEQ ID NO:15:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

10 Gln Gln His Asn Glu Tyr Pro Leu Thr
1 5 9

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
15 (B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Pro Lys Asn Ser Ser Met Ile Ser Asn Thr Pro
1 5 10 11

20 (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 7 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

His Gln Ser Leu Gly Thr Gln
1 5 7

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
30 (A) LENGTH: 8 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

35 His Gln Asn Leu Ser Asp Gly Lys
1 5 8

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: Amino Acid

(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

His Gln Asn Ile Ser Asp Gly Lys
1 5 8

5 (2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Val Ile Ser Ser His Leu Gly Gln
1 5 8

(2) INFORMATION FOR SEQ ID NO:21:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

20 CACTTTGGAT ACCGCGTCCT GCAGGT 26

(2) INFORMATION FOR SEQ ID NO:22:

25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CATCCTGCAG GTCTGCCTTC AGGTCA 26

(2) INFORMATION FOR SEQ ID NO:23:

30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

35 Met Ile Ala Pro Ala Ser Ser Ser Thr Arg Tyr Asn Gln Lys Phe
1 5 10 15

Lys Asp

17

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 121 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	
	1				5					10					15	
10	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	
					20					25					30	
	Gly	His	Trp	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	
					35					40					45	
15	Glu	Trp	Val	Gly	Met	Ile	Ala	Pro	Ala	Ser	Ser	Ser	Thr	Arg	Tyr	
					50					55					60	
	Asn	Gln	Lys	Phe	Lys	Asp	Arg	Phe	Thr	Ile	Ser	Val	Asp	Lys	Ser	
					65					70					75	
	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	
					80					85					90	
20	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Gly	Ile	Tyr	Phe	Tyr	Gly	Thr	
					95					100					105	
	Thr	Tyr	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	
					110					115					120	
25	Ser															
	121															

WHAT IS CLAIMED IS:

1. A humanized anti-CD11a antibody which binds specifically to human CD11a I-domain.
2. The humanized anti-CD11a antibody of claim 1 which binds to epitope MHM24 on CD11a.
3. A humanized anti-CD11a antibody which binds human CD11a with a K_d value of no more
5 than about 1×10^{-8} M.
4. A humanized anti-CD11a antibody which has an $IC_{50}(nM)$ value of no more than about 1nM
for preventing adhesion of Jurkat cells to normal human epidermal keratinocytes expressing ICAM-1.
5. A humanized anti-CD11a antibody which has an IC_{50} (nM) value of no more than about
1nM in the mixed lymphocyte response assay.
- 10 6. The humanized anti-CD11a antibody of claim 1 having a heavy chain variable region
comprising the amino acid sequence of CDR1 (SEQ ID NO:10), CDR2 (SEQ ID NO:11) and CDR3 (SEQ
ID NO:12) of humanized antibody MHM24 (F(ab)-8.
7. The humanized anti-CD11a antibody of claim 6 comprising the amino acid sequence of SEQ
ID NO:5.
- 15 8. The humanized anti-CD11a antibody of claim 1 having a light chain variable region
comprising the amino acid sequence of CDR1 (SEQ ID NO:13), CDR2 (SEQ ID NO:14) and CDR3 (SEQ
ID NO:15) of humanized MHM24 F(ab)-8.
9. The humanized anti-CD11a antibody of claim 8 comprising the amino acid sequence of SEQ
ID NO:2.
- 20 10. The humanized anti-CD11a antibody of claim 1 having a light chain variable region
comprising the amino acid sequence of SEQ ID NO:5 and a light chain variable region comprising the amino
acid sequence of SEQ ID NO:2.
11. The humanized anti-CD11a antibody of claim 1 which is a full length antibody.
12. The humanized anti-CD11a antibody of claim 11 which is a human IgG.
- 25 13. The antibody of claim 1 which is an antibody fragment.
14. The antibody fragment of claim 13 which is a $F(ab')_2$.
15. The antibody of claim 1 which is labeled with a detectable label.
16. The antibody of claim 1 which is immobilized on a solid phase.
17. The antibody of claim 1 which is conjugated to a cytotoxic agent.
- 30 18. A method for determining the presence of CD11a protein comprising exposing a sample
suspected of containing the CD11a protein to the antibody of claim 1 and determining binding of said antibody
to the sample.
19. A kit comprising the antibody of claim 1 and instructions for using the antibody to detect
the CD11a protein.
- 35 20. Isolated nucleic acid encoding the antibody of claim 1.
21. A vector comprising the nucleic acid of claim 20.
22. A host cell comprising the vector of claim 21.
23. A process of producing a humanized anti-CD11a antibody comprising culturing the host cell
of claim 22 so that the nucleic acid is expressed.

24. The process of claim 23 further comprising recovering the humanized anti-CD11a antibody from the host cell culture.

25. The process of claim 24 wherein the humanized anti-CD11a antibody is recovered from the host cell culture medium.

5

1 / 4



	1	10	20	30	40
MHM24	DVQITQSPSYLAASPGETISINC [RASKTISKYLA] WYQEKPGKTNKLLIY				
				
F(ab)-8	DIQMTQSPSSLSASVGDRVITTC [RASKTISKYLA] WYQQKPGKAPKLLIY				
				
humKI	DIQMTQSPSSLSASVGDRVITTC [RASQISNYLA] WYQQKPGKAPKLLIY				
	60	70	80	90	
MHM24	[SGSTLQS] GIPSRFSGSGSGTDFTLTISLQPEDFAMYIC [QQHNEYPLT]				
				
F(ab)-8	[SGSTLQS] GVPSRFSGSGSGTDFTLTISLQPEDFATYYC [QQHNEYPLT]				
				
humKI	[AASSLES] GVPSRFSGSGSGTDFTLTISLQPEDFATYYC [QQYNSLPWT]				
	100				
MHM24	FGTGTKLELKR				
	. . .				
F(ab)-8	FGQGTKVEIKR				
	. . .				
humKI	FGQGTKVEIKR				

FIG. 1A



	1	10	20	30	40		
MHM24	EVQLQQPGAELMRPGASVKLSCKASGYSFT [GHWMN] WVRQRPQGGLWIG						
						
F(ab)-8	EVQLVESGGGLVQPGGSLRLSCAASGYSFT [GHWMN] WVRQAPGKGLEWVG						
						
humIII	EVQLVESGGGLVQPGGSLRLSCAASGFTFS [SYAMS] WVRQAPGKGLEWVS						
RhIgG1	EVQLVESGGGLVQPGGSLRLSCAASGYSFT [GHWMN] WVRQAPGKGLEWVG						
	50	a	60	70	80	abc	90
MHM24	[MIHPSDSETRLNQKFKD] KATLTVDKSSSSAYMQLSSPTSSED SAVYYCAR						
						
F(ab)-8	[MIHPSDSETRYNQKFKD] RFTISVDKSKNTLYLQMNSLRAEDTAVYYCAR						
						
humIII	[VISGDGGSTYADSVKG] RFTISRDNSKNTLYLQMNSLRAEDTAVYYCAR						
RhIgG1	[MIA P ASSSTRYNQKFKD] RFTISVDKSKNTLYLQMNSLRAEDTAVYYCAR						
	100		110				
MHM24	[GIYFYGTTYFDY] WGQGTTLTVSS						
						
F(ab)-8	[GIYFYGTTYFDY] WGQGTTLTVSS						
						
humIII	[G-----FDY] WGQGTTLTVSS						
RhIgG1	[GIYFYGTTYFDY] WGQGTTLTVSS						

FIG. 1B



2 / 4

**FIG. 2****FIG. 3**

3 / 4



4 / 4

**FIG._4C****FIG._5**

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/19041

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/62 C07K16/28 C07K16/46

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	J. HILDRETH ET AL.: "A human lymphocyte-associated antigen involved in cell-mediated lympholysis." EUROPEAN JOURNAL OF IMMUNOLOGY, vol. 13, no. 3, March 1983, WEINHEIM, GERMANY, pages 202-208, XP002054055 cited in the application see the whole document --- -/-	1-25

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

30 January 1998

Date of mailing of the international search report

20/02/1998

Name and mailing address of the ISA

European Patent Office. P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Nooij, F

INTERNATIONAL SEARCH REPORT

Intern 1st Application No

PCT/US 97/19041

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	C. EDWARDS ET AL.: "Identification of amino acids in the CD11a I-domain important for binding of the leukocyte function-associated antigen (LFA-1) to intercellular adhesion molecule-1 (ICAM-1)." THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 21, 26 May 1995, BALTIMORE, MD, USA, pages 12635-12640, XP002054056 see abstract see figure 1 ---	1-25
A	WO 96 32478 A (GENENTECH, INC.) 17 October 1996 see claims ---	1-25
A	EP 0 440 351 A (MERCK & CO. INC.) 7 August 1991 see claims ---	1-25
A	P. CARTER ET AL.: "Humanization of an anti-p185HER2 antibody for human cancer therapy." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 89, no. 10, May 1992, WASHINGTON, DC, USA, pages 4285-4289, XP000275844 cited in the application see the whole document ---	1-25
P,X	W. WERTHER ET AL.: "Humanization of an anti-lymphocyte function-associated antigen (LFA-1) monoclonal antibody and reengineering of the humanized antibody for binding to rhesus LFA-1." THE JOURNAL OF IMMUNOLOGY, vol. 157, no. 11, 1 December 1996, BALTIMORE, MD, USA, pages 4986-4995, XP002054057 see the whole document -----	1-25

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/19041

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9632478 A	17-10-96	AU 5435996 A	30-10-96
EP 440351 A	07-08-91	AU 635996 B	08-04-93
		AU 6984391 A	25-07-91
		CA 2034574 A	20-07-91
		CA 2034593 A	20-07-91
		EP 0438312 A	24-07-91
		JP 6054697 A	01-03-94
		JP 6054698 A	01-03-94